

Understanding the potential of urban parks to promote well-being

Received: 27 August 2024

Accepted: 9 October 2025

Published online: 21 November 2025



Linus W. Dietz¹✉, Sanja Šćepanović^{2,3}, Ke Zhou^{2,4}, André Felipe Zanella⁵ & Daniele Quercia^{1,2,6}

Urban parks are important for public health, but the role of specific spaces, such as playgrounds or lakes, and elements, such as benches or sports equipment, in supporting well-being is not well understood. Here, based on expert input and a review of the literature, we defined six types of health-related activity: physical, mindfulness, nature appreciation, environmental, social and cultural. We built a lexicon that links each activity to specific elements and spaces within parks present in OpenStreetMap. Using these data, we scored 23,477 parks across 35 cities worldwide on the basis of their ability to support these activities. We found clear patterns: parks in North America focus more on physical activity, while those in Europe offer more chances to enjoy nature. Parks near city centers support health-promoting activities better than those farther out. Suburban parks in many cities lack the spaces and equipment needed for nature-based, social and cultural activities. We also found large gaps in park quality between cities. Tokyo and Paris provide more equal access, while Copenhagen and Rio de Janeiro show sharp contrasts. These results can help cities create fairer parks that better support public health.

As the world's population continues to gravitate toward urban areas, cities are faced with the immense task of creating and maintaining green spaces to foster public health^{1,2}. Urban parks are especially beneficial to vulnerable population groups such as socioeconomically deprived³ and older people^{4,5}. Yet, the provision of amenities and facilities that support health-promoting activities has not been systematically analyzed at scale.

Urban green spaces can support health in five main ways. First, they help people stay active, which can reduce obesity and heart diseases^{6,7}. Second, they give people a calm space to rest, reduce stress and improve focus^{6,8–10}. Third, they bring people together and support social ties¹¹. Fourth, they clean the air, block noise and cool the city^{6,12}. Last, some green areas expose people to biodiverse forms of life, which may improve the immune system^{13,14}.

Park planners have often focused on practical ways to boost public health, such as adding sports fields, paths or playgrounds^{15–18}. Despite growing evidence of positive effects, spaces that allowed a diverse

range of people to relax or enjoy beauty were often neglected in the past¹⁹. Parks also have the power to build stronger communities²⁰, and many planners now recognize that parks offer clear environmental gains such as cleaner air and lower noise levels¹². During the coronavirus disease 2019 (COVID-19) pandemic, cities saw how vital parks are for public well-being. This led to stronger calls for fair and flexible park design²¹.

Most past studies have treated parks as simple green zones, based on size or distance from homes^{22,23}. Fewer have studied how the park layout and equipment affect how people use them. For instance, SOPARC is a tool that tracks how people move through parks and what spaces they use²⁴. Other tools look at what makes parks good for activity, such as access, safety or looks²⁵. Still others focus on blue spaces such as rivers or lakes²⁶. These tools work well but are time-consuming to implement and hard to scale, and most focus only on physical activity^{27,28}. We still lack a clear understanding of the net benefits of parks, especially when weighing their positive aspects, such as providing spaces for recreation

¹Department of Informatics, King's College London, London, UK. ²Social & Responsible AI Research, Nokia Bell Labs, Cambridge, UK. ³The Uehiro Oxford Institute, University of Oxford, Oxford, UK. ⁴School of Computer Science, University of Nottingham, Nottingham, UK. ⁵Telefónica Innovación Digital, Barcelona, Spain. ⁶Department of Control and Computer Engineering, Politecnico di Torino, Turin, Italy. ✉e-mail: linus.dietz@kcl.ac.uk




Fig. 1 | Scores for nature appreciation and physical activities in London parks. **a,b**, Scores for nature appreciation (**a**) and physical activities (**b**) in London parks. Some parks score low on nature appreciation but high on physical activity (A), whereas imperial-era parks, such as Green Park and St. James's Park near the royal palace, score high on nature appreciation but low on physical activity (B).

Queen Elizabeth Olympic Park, purpose-built for the 2012 Games, scores high across all dimensions (C). A visualization for all cities can be viewed at <https://social-dynamics.net/healthy-parks>. Map data from OpenStreetMap under the Open Data Commons Open Database License (ODbL).

and supporting urban health^{3,29}, with potential downsides, including reinforcing disparities^{30–32} or contributing to gentrification^{33,34}.

For many cities, there is no easy way to list what parks contain or link these parts to health uses. In this study, we score parks in 35 cities on the basis of how well they support six types of health-related activity. The project page is available at <https://social-dynamics.net/healthy-parks>. We then ask how these scores relate to fair access to well equipped parks. Our work answers three questions. First, which health-related activities do different park spaces support? Second, how can we measure parks around the world based on this support? Third, where are the biggest gaps in what parks provide?

Results

We began by creating a lexicon that linked park spaces and elements to health-related activities using OpenStreetMap (OSM) data. We then scored parks in 35 major cities, which we selected for their geographic diversity and reliable OSM coverage. After validating these scores through statistical comparison and expert review, we analyzed differences in park offerings within and between cities.

Lexicon of health-related park features

Defining activities in parks (step 1). An expert panel of three researchers with expertise in urban computing, Earth observation and computational social science, identified common park activities and organized them into six categories:

- Physical activities: movement and sport such as walking, biking, swimming and group exercise;
- Mindfulness activities: practices such as yoga, meditation and tai chi;
- Nature appreciation: observing and enjoying the natural environment, including bird watching and picnicking;
- Environmental activities: community involvement such as gardening and conservation;
- Social activities: gatherings and group events such as festivals and volunteering;
- Cultural activities: heritage and arts programs, including performances and exhibitions.

While our six categories occasionally overlapped, they captured park activities at a more useful granular level than the typical recreational and physical division. Some overlap is inevitable because certain activities such as football (with social and physical aspects) are inherently multifaceted.

Linking facilities to activities (step 2). We used OSM to collect elements and spaces located within parks in the 35 cities. These included defined areas such as forests and ponds as well as features such as benches and sports courts. Each item was described with a tag, which we used to assign the item to an activity category. To scale this process, we trained a classifier based on a large language model (LLM) and validated it with an expert-coded dataset. The full lexicon included 1,441 OSM tags. The ten most frequent tags for each category are listed in Extended Data Table 1. No OSM tags matched the mindfulness category, so we did not include it in our scoring. Future research may incorporate behavioral data to address this gap.

Scoring parks for health support

We scored each park based on the number of activity-related elements and spaces it contained, normalized by park area. Each park received a score for every activity category. Figure 1 presents an example of these scores for London. In total, we scored 23,477 parks across 35 cities on 5 continents (Table 1). Each score reflects how well a park supports a specific activity compared with other parks of similar size in the same city. Figure 2 visualizes the methodology to estimate the 'average park line'.

Validation with Flickr data. We validated our park scores using geotagged Flickr images taken within park boundaries. These images included both user-generated and machine-generated tags. We used sentence embeddings to match these tags to our activity taxonomy, allowing us to estimate activity scores for each park based on photo content.

We compared these Flickr-based activity scores with our OSM-derived scores using the Pearson correlation coefficient (PCC). Across the 35 cities, the average correlation was 0.39 with a standard deviation of 0.07, indicating moderate but consistent agreement between the two sources. The highest mean (μ) agreement appeared in the cultural ($\mu = 0.53$), social ($\mu = 0.39$) and environmental ($\mu = 0.39$) categories. These types of activity often involve distinctive and photogenic features such as festivals, artworks or gardens, which are likely to be photographed and tagged.

By contrast, the physical ($\mu = 0.30$) and nature appreciation ($\mu = 0.33$) categories showed weaker correlations. We believe this may be because users often photograph people or scenery without tagging specific sports or nature-related elements, making it harder to detect those activities in the data.

At the city level, Washington, DC ($\mu = 0.52$) and Perth ($\mu = 0.50$) had the strongest correlations. These cities are in English-speaking

Table 1 | Our activity scores correlate with the prevalence of photographed activities from Flickr images

City	Parks	Mean PCC	Individual activity categories PCC				
			Physical	Nature appreciation	Environmental	Social	Cultural
Amsterdam	25	0.32	0.24	0.36	0.2	0.49	0.29
Auckland	33	0.46	0.38	0.39	0.43	0.47	0.63
Berlin	77	0.34	0.22	0.14	0.53	0.39	0.42
Boston	50	0.4	0.27	0.43	0.36	0.35	0.58
Buenos Aires	54	0.36	−0.04	0.34	0.92	0.15	0.41
Chicago	69	0.41	0.2	0.45	0.35	0.45	0.6
Christchurch	16	0.49	0.27	0.52	0.21	0.72	0.71
Copenhagen	19	0.33	0.31	0.31	0.01	0.22	0.8
Hong Kong	80	0.3	0.38	0.21	0.36	0.21	0.32
Houston	28	0.46	0.36	0.48	0.15	0.61	0.68
London	304	0.44	0.45	0.4	0.32	0.45	0.55
Madrid	42	0.32	0.08	−0.04	0.19	0.66	0.72
Melbourne	59	0.49	0.45	0.63	0.44	0.64	0.29
Montreal	55	0.43	0.31	0.27	0.44	0.49	0.64
Moscow	65	0.34	0.33	0.2	0.34	0.21	0.6
New York	210	0.42	0.41	0.21	0.58	0.45	0.47
Paris	108	0.39	0.42	0.28	0.3	0.39	0.55
Perth	23	0.5	0.18	0.71	0.3	0.68	0.61
Philadelphia	39	0.46	0.43	0.2	0.63	0.32	0.75
Rio de Janeiro	19	0.36	0.26	0.6	0.23	0.41	0.33
Rome	41	0.31	0.43	0.35	0.08	0.16	0.52
San Diego	47	0.44	0.27	0.38	0.61	0.48	0.44
San Francisco	98	0.31	0.17	0.19	0.42	0.32	0.46
Seattle	76	0.38	0.47	0.36	0.09	0.4	0.6
Seoul	52	0.35	0.34	0.36	0.43	0.23	0.38
Singapore	73	0.36	0.14	0.07	0.61	0.39	0.6
St. Petersburg	28	0.34	0.17	0.22	0.38	0.39	0.56
Stockholm	50	0.47	0.45	0.32	0.93	0.23	0.42
Sydney	99	0.31	0.42	0.39	0.22	0.14	0.36
Taipeh	107	0.34	0.16	0.22	0.48	0.26	0.56
Tokyo	208	0.31	0.19	0.24	0.37	0.32	0.42
Toronto	111	0.34	0.31	0.3	0.17	0.39	0.51
Vancouver	62	0.44	0.33	0.37	0.33	0.5	0.67
Vienna	40	0.3	0.18	0.08	0.57	0.19	0.5
Washington, DC	61	0.52	0.4	0.52	0.56	0.56	0.56
Mean (s.d.)	72.23 (60.07)	0.39 (0.07)	0.29 (0.12)	0.33 (0.16)	0.39 (0.21)	0.39 (0.16)	0.53 (0.13)

PCCs between OSM scores and Flickr activity scores for parks with at least 250 photos.

countries where Flickr and OSM usage is more common, which probably led to higher data quality and tag relevance. Cities with lower correlations, including Amsterdam, Hong Kong and Vienna (each $\mu \approx 0.3$), may have had fewer geotagged images, lower tag accuracy or less alignment with our English-based taxonomy.

These findings support the use of OSM data to assess park infrastructure for health-promoting activities, especially in categories and regions where online content about park use is widely available.

Validation with Wikipedia. We also validated our activity scores using Wikipedia pages. We identified the top-scoring parks globally for each health-related activity based on our scoring system. For each park, we

examined the Wikipedia page to confirm whether the park was known for the corresponding activity type.

In the physical category, we found that parks such as Centennial Parklands in Australia and Bois de Boulogne in Paris were known for extensive sports infrastructure, including fields for tennis, soccer and polo. For nature appreciation, top parks such as La Dehesa de la Villa in Madrid and Guandu Nature Park in Taipei stood out in their cities with large green areas and wildlife. In the environmental category, Washington Park Arboretum in Seattle and Kita-no-maru Park in Tokyo included botanical gardens and conservation efforts. Social activity leaders such as Inspiration Lake in Hong Kong and Toronto Island Park in Canada were cited for hosting gatherings and public amenities.




Fig. 2 | Visualization of the scoring method. a,b, The linear regression models for determining the park health scores for the park elements (a) and park spaces (b) in London, UK. The horizontal axis denotes the park's area ($\log_2$), while the

vertical axis represents the count of park elements ($\log_2$) and the area occupied by health-promoting park spaces ($\log_2$). The modest R^2 values are anticipated, highlighting the variety among parks.

In the cultural category, Ueno Park in Tokyo featured museums and historic landmarks, while Seattle Center included concert halls and art venues.

To further verify our assignments, we conducted structured search queries, such as 'Tokyo parks for cultural activities' and found that the top-ranked parks by our method consistently appeared in online recommendations and guides. A complete list of these parks, organized by city and activity category, is available in Extended Data Table 2.

Cities differ in health-related activity support

Our analysis revealed systematic variations in how urban parks support health-promoting activities, with differences occurring both between cities and within individual urban areas. These variations reflect distinct urban planning priorities, geographic influences and equity considerations that shape park design and resource allocation.

Estimating park offerings through 8-hectare model parks. To compare park features systematically across cities, we constructed statistical models of typical 8-hectare parks: a size representing neighborhood parks and matching our dataset's average. Figure 3 (top) shows the expected number of elements and area dedicated to each activity type in a statistically average 8-hectare park for each city. We computed these values using parameters from linear regression models developed for each city (equation (1)). While we fit separate models per city, making raw health scores not directly comparable between cities, the models enabled comparison of hypothetical average parks across urban contexts.

European nature focus versus North American physical activity in regional priorities. The most notable regional contrast lies in emphasis on physical activity versus nature appreciation, as shown in Fig. 3 (bottom). Parks in European cities tended to provide more space for nature appreciation, while North American cities allocated more area to physical activities. These differences reflect broader urban planning

patterns and suggest that parks serve different roles depending on regional goals and cultural expectations.

Universally limited support for cultural features. Across all regions, cultural features appeared least frequently in parks. In terms of elements, nature appreciation appeared most often, followed by social and environmental activities. Cultural features were the least common. This pattern suggests that, while nature and movement receive widespread support, cultural health-promoting activities are deprioritized globally in urban park design.

Park offerings are unequally distributed within cities

We examined how park scores vary within cities by grouping parks into quartiles based on distance from city centers, from the innermost 25% (Q1) to the outermost 25% (Q4). This approach allowed us to systematically assess whether proximity to urban centers influences park offerings. Extended Data Fig. 1 visualizes the negative correlations between park health scores and the distance to the city centers.

Inner-city parks consistently outperform peripheral ones. Across all activity types, parks in central areas scored higher. Figure 4a demonstrates that park scores dropped steadily with each quartile further from the center. This pattern held across most cities on all continents, with exceptions in Buenos Aires and Rio de Janeiro, where geography and coastal form influence green space layout.

Nature appreciation shows strongest distance-related decline. The trend was most pronounced for nature appreciation. This finding challenges assumptions that outer areas naturally support greener, more restorative environments and echo Montgomery's concept of the 'Savannah Trap'³⁵, where suburban areas include empty, open land that does not support social or ecological use.

We systematically verified that the trends in Fig. 4a are statistically significant, as shown in Extended Data Table 3.




Fig. 3 | Area of park spaces and count of elements in a typical 8-hectare park by activity and city. a, Area of park spaces (horizontal axis) and count of elements (vertical axis) in a typical 8-hectare park by activity and city. Nature and physical activity spaces are the most frequent. Nature-related elements are also the most common across cities. Cultural spaces and elements appear least often. Axes are

scaled for clarity. **b**, Average space in 8-hectare parks dedicated to nature and physical activities in North America ($n = 24$) and Europe ($n = 22$). North American parks support physical activity more, while European parks provide more space for nature appreciation. Standard box plot with Q1, median, Q3; whiskers at 1.5 $\times$ interquartile range.

Cities vary in ensuring equitable park access

We assessed equity in park offerings within cities using a disparity index (equation (3)) ranging from 0 (complete equality) to 1 (maximum inequality). Box plots in Fig. 4b,c show average disparities by activity and region, with detailed values in Table 2. Extended Data Fig. 2 visualizes the disparity scores subdivided by city.

Activity-specific disparity patterns. Disparities varied by activity type: physical activities showed the most even distribution ($\mu = 0.135$; $\sigma = 0.039$), while social activities exhibited the highest disparity ($\mu = 0.222$, $\sigma = 0.073$), indicating that some parks offer rich social features while others lack them entirely. Nature appreciation had moderate average disparity ($\mu = 0.157$) but low variation ($\sigma = 0.032$), suggesting consistent global trends. Overall disparities averaged 0.165, and were similar across continents (0.15–0.19), suggesting that inequality in park design is not limited to one region.

Cities achieving balanced activity support. Tokyo, Paris, Auckland, Buenos Aires and Hong Kong demonstrated the most equitable park offerings across activities. Tokyo and Paris particularly excelled, with four or more activity categories showing disparities below one standard deviation from the mean. These results may reflect centralized park governance, strong public policy and active community involvement in cities such as Paris³⁶ and Tokyo³⁷.

Cities with concentrated park resources. By contrast, Moscow, St. Petersburg, Stockholm, Rio de Janeiro and Copenhagen showed high disparities in park offerings across categories. Stockholm exhibited the highest single-category disparity in cultural features (0.514), suggesting cultural resources were concentrated in a few parks. Copenhagen showed high disparity in both cultural and social categories,

potentially reflecting gentrification³⁸. Rio de Janeiro displayed widespread disparities across social, environmental and cultural categories, aligning with prior findings on spatial health inequality³⁹.

These findings demonstrate that, while park inequality exists globally, specific cities have successfully implemented more equitable approaches to supporting diverse health-promoting activities through their urban park systems.

Discussion

Urban parks offer health benefits that go beyond the usual physical and mental well-being. However, our global analysis reveals that urban parks require strategic improvement focused on geographic equity and activity diversity to maximize their health benefits.

Main findings

By identifying six types of activity people do in parks, we created a detailed taxonomy of park activities and used this to evaluate parks worldwide based on their support for five of the six activity categories (data to evaluate the mindfulness category were not available on OSM). By giving individual scores for each activity, we identified three critical areas that demand attention from landscape architects, urban designers and policymakers.

Geographic inequities demand immediate action. Our findings show a clear and concerning geographic trend: parks in city centers are better equipped for health-promoting activities than those on the outskirts. This pattern holds across all types of activity, including those that might be expected to be more common in less populated areas, such as nature appreciation or environmental activities. This geographic disparity contradicts basic expectations and creates notable barriers to equitable health access.




Fig. 4 | Inequalities observed in the spatial distribution of parks and across activity categories. **a**, Park health scores by quartiles of the distance to the city center. Q1 includes parks closest to the city center; Q4 includes those farthest away. Scores decline steadily with distance, especially for nature appreciation. Environmental features show the least decline. Standard box plot with Q1, median, Q3; whiskers at 1.5× interquartile range. $n = 5,869$ parks per quartile.

b,c, Inequality in park health scores activity category (**b**, $n = 35$ cities) and continent (**c**, $n =$ cities per continent; Table 1). While total disparity ranges are similar across regions, some activities, such as social and cultural, show much higher inequality than others. White × indicates mean value. Standard box plot with Q1, median, Q3; whiskers at 1.5× interquartile range.

This finding aligns with previous research. Wolch et al. found that park distribution often favors more wealthy communities³¹. Other studies have shown that recent urban development policies have led to substantial greening in city centers, while suburban areas have received less investment⁴⁰. Our results expand on these works by showing not only reduced access to parks in suburban areas^{31,40}, but also a lack of amenities and spaces for health-promoting activities.

This evidence creates a clear call to action for urban planners to shift their focus to improving suburban parks, as well as parks outside urban centers, where notable and often overlooked recreation gaps are often found⁴¹. However, creating high-quality urban green spaces can be a balancing act, with potential downsides such as gentrification^{31,34}. Designers must think carefully about the urban context when improving parks in ways that may change their role³⁴.

Activity-type inequities show mixed but encouraging patterns. We also examined disparities in the health potential of parks among cities by activity type. The findings are generally encouraging: overall disparity scores were low and did not show pronounced global variation, in contrast to earlier studies^{42,43}. While previous work has shown high inequality in absolute access to greenery, our results suggest that relative inequality in access to health-promoting park activities is less severe.

The distribution patterns vary considerably by activity type, revealing both strengths and critical gaps. Physical activity infrastructure was the most evenly distributed, probably reflecting its prominence in public health discourse and urban policy¹⁸. This aligns with the medical literature's emphasis on physical activity benefits. By contrast, facilities for social activities were the most unequally distributed, revealing a key area for improvement⁴⁴.

The global nature of this challenge is evident from our data: the five cities with the greatest disparities in park offerings span four continents, indicating that uneven provision of well-equipped parks is not a concentrated issue. This suggests that urban planning is not only a matter of financial resources but also of municipal priorities and community engagement^{36,45}. A positive trend can be observed in the Asian countries of our study and Oceania, which have the most fair provision of park facilities.

Our findings provide actionable insights for the design of well-equipped parks. To understand the relevance of our work for urban planning, we conducted 30-min semi-structured interviews with three domain experts. We reached out to practitioners with 12–30 years of professional experience from diverse backgrounds: a lecturer and urban designer (E1, male), an urban designer and master planner (E2, female) and a municipal park development manager (E3, male). The experts confirmed that our quantitative approach offers significant practical value for urban design decisions. They identified several ways our findings could support urban design: reducing reliance on subjective park assessments (E1), serving as a quantitative baseline for master planning (E1 and E2) and providing an evidence-based alternative to institutional knowledge and practice-specific heuristics (E1). The urban designers emphasized that the visualization of park scores is already useful to "identify the gaps regarding what is present in terms of offerings in a district (E1)", especially when aiming to ensure fair access to parks. This was echoed by the park manager, who stated they would use the park scoring to guide development priorities, and as supporting evidence to gain leadership backing or external funding. When shown the taxonomy of activities, they noted that most of their projects addressed physical and nature-appreciation activities, and

Table 2 | The disparity scores by city and activity type are generally low, but vary based on the activity category

City	Mean score	Inequality of each activity category's offering in a city.				
		Physical	Nature appreciation	Environmental	Social	Cultural
Tokyo	0.089	0.065	0.082	0.066	0.146	0.084
Paris	0.097	0.104	0.101	0.087	0.114	0.079
Auckland	0.121	0.103	0.117	0.105	0.154	0.126
Buenos Aires	0.125	0.172	0.107	0.098	0.137	0.108
Hong Kong	0.125	0.079	0.134	0.197	0.094	0.12
Vienna	0.135	0.13	0.144	0.107	0.17	0.125
New York	0.138	0.113	0.137	0.131	0.177	0.129
Rome	0.138	0.122	0.16	0.116	0.153	0.141
Christchurch	0.138	0.104	0.113	0.101	0.273	0.101
San Francisco	0.14	0.09	0.152	0.116	0.212	0.132
Berlin	0.143	0.131	0.14	0.138	0.152	0.154
Chicago	0.153	0.077	0.178	0.205	0.151	0.153
San Diego	0.153	0.129	0.169	0.124	0.214	0.128
Melbourne	0.155	0.122	0.167	0.124	0.248	0.114
Montreal	0.156	0.126	0.178	0.142	0.215	0.118
Singapore	0.157	0.166	0.14	0.119	0.227	0.133
Perth	0.157	0.132	0.19	0.125	0.21	0.128
Madrid	0.16	0.104	0.163	0.168	0.197	0.167
Sydney	0.16	0.11	0.119	0.153	0.226	0.191
London	0.163	0.134	0.154	0.138	0.23	0.158
Vancouver	0.165	0.102	0.144	0.153	0.216	0.208
Toronto	0.169	0.129	0.178	0.124	0.24	0.172
Houston	0.173	0.139	0.188	0.141	0.218	0.18
Taipeh	0.174	0.082	0.156	0.116	0.138	0.377
Philadelphia	0.175	0.158	0.174	0.122	0.308	0.116
Seattle	0.179	0.132	0.161	0.141	0.265	0.195
Boston	0.186	0.158	0.183	0.147	0.292	0.152
Washington, DC	0.191	0.154	0.16	0.173	0.262	0.205
Seoul	0.191	0.177	0.184	0.174	0.271	0.152
Amsterdam	0.197	0.161	0.187	0.153	0.267	0.217
Moscow	0.205	0.232	0.161	0.205	0.236	0.194
St. Petersburg	0.213	0.21	0.166	0.288	0.201	0.199
Stockholm	0.247	0.189	0.132	0.105	0.297	0.514
Rio De Janeiro	0.254	0.141	0.193	0.245	0.455	0.236
Copenhagen	0.266	0.209	0.209	0.195	0.398	0.321
Mean (s.d.)	0.165 (0.039)	0.134 (0.039)	0.155 (0.029)	0.144 (0.045)	0.222 (0.074)	0.172 (0.085)

Cities are ordered by their overall average disparity. Lower values indicate a more even distribution of features. Scores one standard deviation below or above the mean are marked in italics and bold, respectively.

"[the activity taxonomy] helps us think about what it is we're doing and how and how some categories have been neglected (E3)".

The experts also highlighted the importance of the way our scores are normalized to the specific context of each city. Our park scores are normalized to reflect the unique context of each city, making direct comparisons between cities hard. While this limits intercity comparisons, it avoids drawing misleading conclusions across culturally, geographically, and climatically diverse settings. For example, E1 cited culturally different approaches to park provision in the form of 'three-dimensional parks' in high-density environments such as Hong Kong, while E2 brought up the challenges in adopting urban transformation manuals from Western cities in regions with extreme climates such as the Middle East.

Limitations and future directions

Despite the well-known benefits of parks, lack of fair access to urban green spaces remains a common theme in academic literature^{2,31,40,42,46}. We focused only on urban parks, excluding other types of urban green space such as gardens, street trees and green roofs, which also contribute to urban health. Parks were selected because they support a wider range of activities than more specialized spaces, are generally publicly accessible and are typically maintained by municipal administrations, making any identified shortcomings more actionable for policy and planning.

Relying on map data limits our analysis to activities that can be linked to specific elements and spaces within parks. While some studies

explored the relationship between open spaces and their use⁴⁷, we avoided making assumptions about the usage of informal spaces to reduce the risk of cultural bias in our park evaluations. As a result, we excluded mindfulness activities from our main analysis, as they often do not require designated areas in city parks. Likewise, our data do not account for temporary cultural events, such as music festivals held in parks.

Park offerings were scored under the assumption that the presence of facilities and spaces enabling certain activities is a necessary condition for realizing specific health benefits. However, their presence alone may not be enough to deliver these benefits in practice. Furthermore, not all activities are directly tied to specific infrastructure, and OSM cartographic data alone cannot capture the intensity of use. We did not distinguish between the quality of maintenance or design of individual elements, such as benches being equipped with backrests, or the size, layout and species of trees and other plants.

Like all urban spaces, parks are in constant flux, which raises the question of how parks have evolved over time². This became clear during the COVID-19 pandemic, when many cities had to rethink their strategies, possibly leading to big changes in park facilities with major social justice effects⁴⁸. Looking at historical data from OSM is challenging, as it is difficult to distinguish between actual changes in park facilities and the increasing completeness of the OSM database.

A key direction for future research is to examine how park offerings directly influence health outcomes. While collecting global medical data to evaluate the health impact of individual parks is impractical, more targeted investigations may be feasible in specific contexts. For instance, prescription data available in some countries could be used to estimate the prevalence of certain medical conditions⁴⁹, and to explore potential causal links between changes in park infrastructure and improvements in public health over time.

We showed that the OSM park scores are closely linked to activities captured in Flickr photos, but other data could also be used to approximate park usage. For instance, detailed mobile traffic data could offer interesting insights into how parks are used^{30,50,51}. This could also help to reduce biases inherent in social media data sources such as Flickr and OSM, which rely on actively submitted contributions from tech-savvy users, and sometimes out-of-city visitors rather than passive observations of citizen behavior.

Finally, by focusing on health-promoting activities, our work introduces avenues for health-related behavioral change^{20,52}. For instance, our activity-based approach could support context-aware recommendations⁵³, such as identifying the most suitable times to engage in specific park activities when conditions such as temperature, air quality or crowdedness are most favorable.

Methods

Data

OSM is a globally encompassing geographic information database based on crowdsourced contributions. While accessible through a map interface at <https://openstreetmap.org>, its primary value lies in being an indispensable source for open mapping data in both commercial and scientific applications^{54–56}. Thanks to its permissive licensing, OSM has fostered a large ecosystem of individual and professional contributors. As a result, the project has achieved comprehensive worldwide coverage, with near-perfect mapping quality across the Western world⁵⁶, while retaining remarkable detail in the global south⁵⁵. In this study, we utilized OSM data from fall 2023.

OSM uses a tagging system with key–value pairs to categorize and describe all these map objects. Each map object is typically associated with multiple tags that describe its purpose, but may also include additional information, such as opening hours, or data source references. To avoid the complexities of the OSM data model, for our tasks, it was sufficient to focus on two key map objects related to parks: park elements and park spaces. Park elements are zero-dimensional points

representing objects such as benches, individual trees and statues. Park spaces refer to areas within the parks, such as meadows, lakes and forests.

Flickr (<https://flickr.com>) has established itself as one of the most prominent platforms for sharing photography. Since its inception in 2004, the platform has gained considerable popularity, accumulating billions of images. Notably, many of these images have been precisely geolocated, thanks to the utilization of the (phone) camera's GPS module.

We utilized a substantial dataset comprising geolocated images posted between 2004 and 2015. This extensive dataset offered us a valuable secondary perspective on activities taking place within the parks of the world. By intersecting these images with the park outlines from OSM, we identified 10,788,686 pictures captured within the boundaries of parks in our study cities. To extract the depicted content from these images, we used user-assigned tags in conjunction with automatically generated computer vision labels^{57,58}.

Study area

Our research focused on 35 cities listed in Table 1, which we selected using three criteria to make our analysis broad yet robust.

First, we selected major cities worldwide with populations of at least 650,000. This threshold includes many of the largest urban areas, such as major European capitals and other densely populated regions where parks play a vital role in public well-being. To improve representation in Oceania, however, we made an exception for Christchurch, New Zealand, which has a smaller population. This first criterion allowed us to examine parks in cities from various parts of the world, each affected by its own climate, history and cultural background.

Second, we only considered cities in countries where at least 80% of the population has Internet access⁵⁹. This ensured we had enough online data (such as tags on OSM or photos on Flickr) for our study. As there are no detailed global data on Internet use in cities specifically, we used the country's overall access to the Internet as our guide. We decided on this threshold upon our preliminary analyses, finding that, in many cities in Africa and South America, there was not enough digital information for our approach, which relies on social media and collaborative mapping data.

Third, we selected cities where, on average, at least one-eighth of park areas are annotated with health-related tags on OSM. As our analysis relied heavily on OSM data, this criterion ensured a minimum level of information on the platform for our study. We settled on this one-eighth threshold after observing that, below this level, the lack of contributor-added tags limited our ability to extract meaningful information. This primarily excluded cities where most tagging was done predominantly automatically through Earth observation that was not accompanied by manual tagging of OSM contributors. This was mainly the case in China, where nongovernmental mapping is restricted⁶⁰.

Identifying health-promoting activities in urban greenery

We identified and categorized park-based activities using input from an expert panel consisting of three co-authors of this study. We compiled a comprehensive list of activities commonly undertaken in urban parks. To collect relevant papers, we used two specific queries of Google Scholar: '(urban) AND (parks OR greenery) AND usage' and '(activities in urban) AND (parks OR greenery)'. From this process, we retrieved the top 50 scholarly articles for each search phrase, resulting in a total of 91 unique papers. We reviewed each article and collected all activities, resulting in a diverse set of activity descriptions varying in granularity. For example, the literature included both broad terms such as 'leisure activities' or 'recreation', as well as more specific categories such as 'physical' and 'social' activities. We also noted plenty of individual activities such as 'walking', 'performing street theater', 'fishing' and 'playing all kinds of different sports'. Subsequently, we convened to categorize the identified activities, with a particular focus on their

potentially different impact on health and on ensuring a consistent level of specificity across categories. Broad terms such as recreation were deemed too general to be analytically useful, whereas distinctions like physical versus social activities were considered meaningful. This process ultimately yielded six distinct categories: physical, mindfulness, nature appreciation, environmental, social and cultural activities.

Annotating park OSM tags with activities using LLM classifiers

To associate different park elements and spaces with health-promoting activities, we annotated OSM tags describing those elements and spaces with activities. This turned out to be a challenging task. OSM is a collaborative platform with some governance and guidelines (https://wiki.openstreetmap.org/wiki/Map_features) for tagging, but the flexible tagging system offers the crowdsourcing contributors substantial freedom. As a result, the data can be inconsistent and fragmented, necessitating thorough cleaning. Each map object, such as park elements and spaces, can be tagged with an unlimited number of tags, offering in-depth descriptions. As a result, we encountered over 30,000 unique key–value pairs associated with park elements and spaces. As our primary focus is on the core functional aspects of these elements, we conducted a data cleaning step (detailed in the Supplementary Information) to remove irrelevant metadata associated with the map objects. This filtering allowed us to focus exclusively on tags relevant to activity-related features, thereby making the annotation process more pertinent to our study.

Using LLMs for annotation. Even for domain experts, linking these tags unequivocally to health-related activities was difficult. For instance, a bench might relate to socializing, enjoying nature or resting after physical activity. Choosing one activity over another often depended on personal experience, as many tags could plausibly refer to multiple activities. Given the large number of items and the specialized nature of the task, we chose an LLM classifier as an alternative to expert annotation or crowdsourcing.

Using LLMs as classifiers offers several advantages as they provide a more objective and consistent approach to annotation, can handle large volumes of data quickly and do so at relatively low cost. In some cases, LLM classifiers have even outperformed crowdworkers⁶¹, who may themselves rely on machine learning tools to complete tasks⁶². Research also suggests that LLM-based annotations can match the quality of those made by domain experts⁶³. Given these benefits, we established a benchmark in our domain to evaluate the feasibility of using LLMs to map OSM tags to health-promoting activities. Three experts manually annotated the 100 most frequent tags, and final labels were selected by majority vote. We then used this expert-labeled dataset to assess the accuracy of the labels generated by different LLM classifiers. The details of the LLM annotation benchmark can be found in the Supplementary Information. The outcome indicated that GPT-4, set at a temperature of 0.9, yielded the best annotation performance of an F_1 score of 0.77.

Operationalization of the taxonomy. Using the taxonomy with six categories of health-promoting activities, and GPT-4 as the best-performing annotation model, we ran the annotation of OSM tags describing park elements and park spaces. These tags were then labeled with one of the health-promoting activities, or ‘none’ if they did not support a particular activity. In doing so, we established a lexicon of park elements and spaces linked to health-promoting activities (Extended Data Table 1). We had to exclude the mindfulness activities category at this stage, as none of the OSM tags found in parks was primarily associated with it.

Computing park health scores by aggregating OSM tags

The core method to characterize parks in terms of their potential for health-promoting activities is based on counting the respective park elements and spaces. These counts are then combined to give each park

an overall score for each health-related category. This score represents the potential health benefits of each park.

Counting health-promoting elements and spaces in parks. In our process of assigning health-promoting activity scores to each park, we first gathered park elements and spaces within each park using the osmium library. We then assigned health-promoting activities to these elements and spaces based on the lexicon created in the previous step (Extended Data Table 1). We discarded any elements or spaces whose tags did not match an activity category. In a few instances, park elements or spaces could fall into more than one health-promoting activity category. For example, apple trees are annotated in OSM with the tags [natural=tree, produce=apple]. In our lexicon of park elements and spaces, we map natural=tree to the nature appreciation and produce=apple to the environmental category. To account for this overlap, we proportionally assign the element as 50% nature appreciation and 50% environmental. More generally, when multiple tags are matched to different categories, we proportionally count the resource based on the number of matched tags, ensuring that its contributions are accurately accounted for, thus avoiding underestimating secondary activities of multipurpose facilities.

Transforming counts into health scores. After tallying up the park elements and spaces within the park, we measured the overall effect of the park in promoting healthy activities within a city. This score should account for the park’s size and the range of facilities it offers for different activities. Our proposed scoring method is based on the following considerations.

- (1) Amount of health-promoting elements and spaces: The number and area of health-promoting facilities determine how much offering there is for each activity.
- (2) Area of the park: The character of a park depends on the concentration of health-promoting facilities. Larger parks must offer more to obtain a high score.
- (3) Diminishing returns with increased count: We posit that, as the count of these elements and spaces increases, the associated benefits exhibit diminishing returns.
- (4) City-specific normalization of park health scores: The value of a park’s facilities for a certain activity is relative to similar facilities in other parks throughout the city.

To reflect these assumptions into our scoring, we proposed a linear regression model to compute the park health scores shown in equation (1). We used the idea of an ‘average park’ in each city to compute a baseline and used the distance of each park to the average park line, that is, the residual, as a score. The average park baseline was determined by computing separate linear regression models for park elements and spaces in each city, estimating the expected amount of facilities relative to the park area.

$$E_{\text{Act}}(\log_2(\text{count}(\text{Act}))) = i + s \times \log_2(\text{park area}) \mid \text{Act} \in \text{Activity Categories}, \quad (1)$$

where i and s represent the intercept and slope of the regression lines, respectively. To obtain regression models for each activity category and both park elements and park spaces, we utilized the binary logarithm to account for the diminishing returns of an increase in park size. For each city, separate regression models were calculated for each activity category, as well as for park elements and park spaces, which is further detailed in the Supplementary Information.

By analyzing the residuals, we identified parks that exceeded expectations (positive residuals) and those that fell short (negative residuals) in providing health-promoting resources for a given activity. We made the linear model dependent on park area so that the resulting scores for health-promoting elements and spaces reflected their density. To reduce the influence of extremely large parks with high

amounts of facilities, we applied the binary logarithm. We calculated these scores separately for each city, rather than using a single global model, to ensure that the results reflected each city's local context. To illustrate this method, we plotted the linear models and the individual park scores in the log–log space for London, UK (Fig. 2). The regression lines denote expected health scores based on park size. The modest R^2 values showed that the number of health-promoting facilities could not be explained by park size but instead reflected different design priorities and the needs of local citizens. Park scores for park elements and spaces are residuals from this average park line in the model, adjusting for park size when determining health scores.

Combining scores from park elements and spaces. The regression models gave us individual scores for park elements and park spaces for each activity. To verify the impact of combining these scores into one combined health score, we examined the co-occurrences of park elements and spaces and found that they represent orthogonal concepts in practice, as shown in Extended Data Table 4. Based on this finding, we computed a combined score by first normalizing the scores of park elements and spaces using the z-score transformation, considering all parks in a city. This normalization allowed us to standardize the scores, making them comparable despite being on different scales. Then, we linearly combined these z scores by averaging them together to create the overall score for the park according to equation (2).

$$\text{Score}(P_{\text{Act}}) = \frac{z(\text{residual}_{\text{elements}}(P_{\text{Act}})) + z(\text{residual}_{\text{spaces}}(P_{\text{Act}}))}{2}, \quad (2)$$

where P denotes an individual park, Act is one of the activity categories, the residual scores for elements and spaces stem from equation (1), and $z()$ indicates the z-score transformation.

The combined and normalized scores of park elements and spaces represent a comprehensive and unified measure of the park's health-promoting amenities and facilities, accounting for both individual elements and cultivated areas. The combination process accounted for the relative importance of each aspect, leading to a more meaningful overall score that represents how well a park is equipped to support performing health-promoting activities. Because the scores are based on z-score-normalized residuals, a value around 0 indicates average support for a given activity, while a score of ± 1 means the park is 1 standard deviation above or below the city-wide average.

Quantifying disparities of park scores

One goal of our study was to quantify disparities in health benefits offered by different parks within a city. To measure the disparities in the presence of amenities and facilities associated with health-promoting activities within a city, we propose the following disparity index. The metric essentially quantifies the inequality of the park health scores, as, generally, one could expect that good park management would provide for a similar amount of features and facilities in all parks of a city. As the park health scores could be negative, we cannot directly use a standard inequality metric, such as the Gini index, but had to min–max normalize the park score before computing the Gini index (equation (3)).

$$\text{Gini}_{\text{Act}}(\{X'_{\text{Act}} | X_{\text{Act}} : P \in C\}), \quad (3)$$

where X_{Act} is the score of activity category $\text{Act} \in [\text{physical, cultural and so on}]$ of a park P in city C , and

$$X' = \frac{X - X_{\min}}{X_{\max} - X_{\min}} \text{ (min-max normalization),}$$

and the Gini index was computed in a standard way:

$$\text{Gini} = \frac{A}{A + B},$$

where A is the area between the Lorenz curve and the line of perfect equality and B is the area beyond the Lorenz curve⁶⁴.

Semantic matching of Flickr labels and OSM tags

In our validation, we leveraged a global dataset of 10.7 million geotagged Flickr images taken within parks across 35 cities. Each image came with user-generated tags, partially annotated by computer vision algorithms. To semantically match these Flickr labels to OSM tags, we used text embeddings, treating the task as an asymmetric semantic search problem. To overcome language diversity in the Flickr labels, we detected the top three non-English languages per city and translated the labels into English using machine translation models. To further improve embedding quality, we enriched OSM tags with concise definitions from the OSM mapping guidelines. The embeddings were generated using the all-mpnet-base-v2 S-BERT model, and matching was done based on cosine similarity, with a threshold of 0.7 to ensure quality. The methodological details are described in the Supplementary Information.

This process yielded 2,171 Flickr-to-OSM matches, with 1,432 corresponding to health-promoting features. To assess accuracy, three experts reviewed the 20 most frequent label–tag matches for London. We aggregated their responses using majority voting. The experts' annotation agreed with 82% of the matchings, which is highly accurate considering they are based solely on individual tags.

Having assured that the matchings are accurate, we proceeded to profile the parks based on the activities associated with the matched OSM tags, following the same scoring approach as what we used for the OSM park elements and spaces (equation (1)). In our validation, we chose a minimum of 250 images from each park and at least 15 parks in each city. This criterion was established to secure a robust number of images for each park, thus ensuring the accuracy of our analysis. This was a mitigation against potential biases that could have been introduced by individual photographers if a park had only a few images.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

The replication package contains tables of the park health scores in the cities available at <https://social-dynamics.net/healthy-parks/replication>. The original OpenStreetMap data used for scoring the parks are publicly available and can be best obtained from one of the third-party download servers, for example from <https://download.geofabrik.de>. The Flickr dataset for the validation cannot be shared due to the terms of conditions of this dataset.

Code availability

The Python and R code to compute park health scores is available at <https://social-dynamics.net/healthy-parks/replication>.

References

- Kadakia, K. T. & Galea, S. Urbanization and the future of population health. *Milbank Quart.* **101**, 153–175 (2023).
- Wu, S., Chen, B., Webster, C., Xu, B. & Gong, P. Improved human greenspace exposure equality during 21st century urbanization. *Nat. Commun.* **14**, 6460 (2023).
- Maas, J., Verheij, R. A., Groenewegen, P. P., de Vries, S. & Spreeuwenberg, P. Green space, urbanity, and health: how strong is the relation? *J. Epidemiol. Commun. Health* **60**, 587–592 (2006).
- Dzhambov, A. M. & Dimitrova, D. D. Elderly visitors of an urban park, health anxiety and individual awareness of nature experiences. *Urban For. Urban Green.* **13**, 806–813 (2014).

5. Takano, T., Nakamura, K. & Watanabe, M. Urban residential environments and senior citizens' longevity in megacity areas: the importance of walkable green spaces. *J. Epidemiol. Commun. Health* **56**, 913–918 (2002).
6. Hartig, T., Mitchell, R., de Vries, S. & Frumkin, H. Nature and health. *Annu. Rev. Public Health* **35**, 207–228 (2014).
7. Twohig-Bennett, C. & Jones, A. The health benefits of the great outdoors: a systematic review and meta-analysis of greenspace exposure and health outcomes. *Environ. Res.* **166**, 628–637 (2018).
8. Ulrich, R. S. View through a window may influence recovery from surgery. *Science* **224**, 420–421 (1984).
9. Ulrich, R. S. in *Healing Gardens: Therapeutic Benefits and Design Recommendations* (eds Cooper, M. C. & Barnes, M.) 27–86 (John Wiley & Sons, 1999).
10. Berman, M. G., Jonides, J. & Kaplan, S. The cognitive benefits of interacting with nature. *Psychol. Sci.* **19**, 1207–1212 (2008).
11. Jennings, V. & Bamkole, O. The relationship between social cohesion and urban green space: an avenue for health promotion. *Int. J. Environ. Res. Public Health* **16**, 452 (2019).
12. Nieuwenhuijsen, M. J., Khreis, H., Triguero-Mas, M., Gascon, M. & Dadvand, P. Fifty shades of green: pathway to healthy urban living. *Epidemiology* **28**, 63–71 (2017).
13. Rook, G. A. W. Regulation of the immune system by biodiversity from the natural environment: an ecosystem service essential to health. *Proc. Natl Acad. Sci. USA* **110**, 18360–18367 (2013).
14. Hanski, I. et al. Environmental biodiversity, human microbiota, and allergy are interrelated. *Proc. Natl Acad. Sci. USA* **109**, 8334–8339 (2012).
15. Barton, H., Thompson, S., Burgess, S. & Grant, M. *The Routledge Handbook of Planning for Health and Well-being: Shaping a Sustainable and Healthy Future* (Routledge, 2015).
16. Gall, A. R. L., Lemaire, N. & Jabot, F. Lessons learned from co-constructing a guide on healthy urban planning and on integrating health issues into environmental impact assessments conducted on french urban development projects. *Impact Assess. Project Appraisal* **36**, 68–80 (2018).
17. Kaczynski, A. T., Potwarka, L. R. & Saelens, B. E. Association of park size, distance, and features with physical activity in neighborhood parks. *Am. J. Public Health* **98**, 1451–1456 (2008).
18. Pietilä, M. et al. Relationships between exposure to urban green spaces, physical activity and self-rated health. *J. Outdoor Recreation Tourism* **10**, 44–54 (2015).
19. Campbell, K. *Rethinking Open Space, Open Space Provision and Management: A Way Forward* (Scottish Executive Central Research Unit, 2001).
20. Campbell, L. K., Svendsen, E. S., Sonti, N. F. & Johnson, M. L. A social assessment of urban parkland: analyzing park use and meaning to inform management and resilience planning. *Environ. Sci. Policy* **62**, 34–44 (2016).
21. Patwary, M. M. et al. Nature exposure and mental health during the COVID-19 pandemic: a navigation guide systematic review with meta-analysis. *Environ. Pollut.* **356**, 124284 (2024).
22. Brown, G. A theory of urban park geography. *J. Leisure Res.* **40**, 589–607 (2008).
23. Tu, X., Huang, G., Wu, J. & Guo, X. How do travel distance and park size influence urban park visits? *Urban For. Urban Green.* **52**, 126689 (2020).
24. McKenzie, T. L., Cohen, D. A., Sehgal, A., Williamson, S. & Golinelli, D. System for observing play and recreation in communities (SOPARC): reliability and feasibility measures. *J. Phys. Activity Health* **3**, S208–S222 (2006).
25. Bedimo-Rung, A. L., Gustat, J., Tompkins, B. J., Rice, J. & Thomson, J. L. Development of a direct observation instrument to measure environmental characteristics of parks for physical activity. *J. Phys. Activity Health* **3**, S176–S189 (2006).
26. Mishra, H. S. et al. The development of a tool for assessing the environmental qualities of urban blue spaces. *Urban For. Urban Green.* **49**, 126575 (2020).
27. Floyd, M. F., Spengler, J. O., Maddock, J. E., Gobster, P. H. & Suau, L. J. Park-based physical activity in diverse communities of two U.S. cities: an observational study. *Am. J. Prevent. Med.* **34**, 299–305 (2008).
28. Kaczynski, A. T., Stanis, S. A. & Besenyi, G. M. Are park proximity and park features related to park use and park-based physical activity among adults? Variations by multiple socio-demographic characteristics. *Int. J. Behav. Nutr. Phys. Activity* **11**, 146 (2014).
29. Jimenez, M. P. et al. Associations between nature exposure and health: a review of the evidence. *Int. J. Environ. Res. Public Health* **18**, 4790 (2021).
30. Xiao, Y., Wang, D. & Fang, J. Exploring the disparities in park access through mobile phone data: evidence from Shanghai, China. *Landsc. Urban Plan.* **181**, 80–91 (2019).
31. Wolch, J. R., Byrne, J. & Newell, J. P. Urban green space, public health, and environmental justice: the challenge of making cities 'just green enough'. *Landsc. Urban Plan.* **125**, 234–244 (2014).
32. Bolte, A.-M., Kistemann, T., Dehbi, Y. & Kötter, T. (Un)just distribution of visible green spaces? A socio-economic window view analysis on residential buildings: the City of Cologne as case study. *J. Geovis. Spatial Anal.* **9**, 17 (2025).
33. Angelovski, I. et al. Green gentrification in European and North American cities. *Nat. Commun.* **13**, 3816 (2022).
34. Rigolon, A. & Németh, J. Green gentrification or 'just green enough': do park location, size and function affect whether a place gentrifies or not? *Urban Stud.* **57**, 402–420 (2020).
35. Montgomery, C. *Happy City* (Penguin, 2015).
36. Jolé, M. The public of parks: in between observation and action. the example of Paris. *Urbani Izziv* **19**, 169–173 (2008).
37. Fujita, K. & Hill, R. C. Residential Income Inequality in Tokyo and Why it Does Not Translate into Class-based Segregation. in *Residential Segregation in Comparative Perspective* (eds Fujita, K & Maloutas, T.) 37–68 (Routledge, 2016).
38. Post, H. C. Best in the world. *Hans Chr. Post* <https://www.hcpost.dk/portfolio-item/best-in-the-world> (accessed 11 June 2025).
39. Bortz, M. et al. Disaggregating health inequalities within Rio de Janeiro, Brazil, 2002–2010, by applying an urban health inequality index. *Cadernos Saude Publ.* **31**, 107–119 (2015).
40. Sun, L., Chen, J., Li, Q. & Huang, D. Dramatic uneven urbanization of large cities throughout the world in recent decades. *Nat. Commun.* **11**, 5366 (2020).
41. Klein-Hewett, H. Classifying rural parks: a case study in Iowa. *J. Park Recreation Admin.* **42**, 1–18 (2024).
42. Chen, B. et al. Contrasting inequality in human exposure to greenspace between cities of global north and global south. *Nat. Commun.* **13**, 4636 (2022).
43. Long, X., Chen, Y., Zhang, Y. & Zhou, Q. Visualizing green space accessibility for more than 4,000 cities across the globe. *Environ. Plan. B* **49**, 1578–1581 (2022).
44. Zhang, R., Peng, S., Sun, F., Deng, L. & Che, Y. Assessing the social equity of urban parks: an improved index integrating multiple quality dimensions and service accessibility. *Cities* **129**, 103839 (2022).
45. Jones, R. Enticement: the role of community involvement in the management of urban parks. *Manag. Leisure* **7**, 18–32 (2002).
46. Kleinschroth, F. et al. Global disparities in urban green space use during the COVID-19 pandemic from a systematic review. *Nat. Cities* **1**, 136–149 (2024).
47. Goličnik, B. & Ward Thompson, C. Emerging relationships between design and use of urban park spaces. *Landsc. Urban Plan.* **94**, 38–53 (2010).

48. Larson, L. R. et al. Urban park use during the COVID-19 pandemic: are socially vulnerable communities disproportionately impacted? *Front. Sustain. Cities* **3**, 710243 (2021).
49. Šćepanović, S. et al. MedSat: a public health dataset for England featuring medical prescriptions and satellite imagery. *Adv. Neural Inf. Process. Syst.* **36**, 77810–77851 (2024).
50. Heikinheimo, V. et al. Understanding the use of urban green spaces from user-generated geographic information. *Landsc. Urban Plan.* **201**, 103845 (2020).
51. Zanella, A. F. et al. The digital life of Parisian parks: multifunctionality and urban context uncovered by mobile application traffic. Preprint at <https://arxiv.org/abs/2508.15516> (2025).
52. Zhang, L., Tan, P. Y. & Diehl, J. A. A conceptual framework for studying urban green spaces effects on health. *J. Urban Ecol.* **3**, jux015 (2017).
53. Dietz, L. W., Šćepanović, S., Zhou, K. & Quercia, D. Exploratory analysis of recommending urban parks for health-promoting activities. In *18th ACM Conference on Recommender Systems* 1131–1135 (ACM, 2024); <https://doi.org/10.1145/3640457.3691712>
54. Li, J., Qin, H., Wang, J. & Li, J. OpenStreetMap-based autonomous navigation for the four wheel-legged robot via 3D-Lidar and CCD camera. *IEEE Trans. Industr. Electron.* **69**, 2708–2717 (2022).
55. Herfort, B., Lautenbach, S., de Albuquerque, J. P., Anderson, J. & Zipf, A. The evolution of humanitarian mapping within the OpenStreetMap community. *Sci. Rep.* **11**, 3037 (2021).
56. Jacobs, K. T. & Mitchell, S. W. OpenStreetMap quality assessment using unsupervised machine learning methods. *Trans. GIS* **24**, 1280–1298 (2020).
57. Li, L.-J., Socher, R. & Fei-Fei, L. Towards total scene understanding: Classification, annotation and segmentation in an automatic framework. In *Conference on Computer Vision and Pattern Recognition* 2036–2043 (IEEE, 2009); <https://doi.org/10.1109/cvpr.2009.5206718>
58. Thomee, B. et al. YFCC100M: the new data in multimedia research. *Commun. ACM* **59**, 64–73 (2016).
59. *Individuals Using the Internet (% of population)*. (World Telecommunication/ICT Indicators Database, International Telecommunication Union, accessed 9 November 2023); <https://data.worldbank.org/indicator/IT.NET.USER.ZS>
60. Lin, W. Volunteered geographic information constructions in a contested terrain: a case of openstreetmap in china. *Geoforum* **89**, 73–82 (2018).
61. Gilardi, F., Alizadeh, M. & Kubli, M. ChatGPT outperforms crowd workers for text-annotation tasks. *Proc. Natl. Acad. Sci. U.S.A.* **120**, 30 (2023).
62. Veselovsky, V. et al. Prevalence and prevention of large language model use in crowd work. *Commun. ACM* **68**, 3 (2025).
63. Ziems, C. et al. Can large language models transform computational social science? *Comput. Linguist. Assoc. Comput. Linguist.* **50**, 1 (2024).
64. Barrow, M. *Statistics for Economics, Accounting and Business Studies* (Pearson, 2017).

Acknowledgements

We thank K. Firth, S. Law and O. Hayes for their valuable expert input. L.W.D. acknowledges support from the European Union's Horizon 2020 research and innovation program under grant agreement no. 869764 (GoGreenRoutes).

Author contributions

L.W.D. conceived and conducted the experiments and codrafted the paper with S.Š. K.Z., A.F.Z. and D.Q. conceived the experiments and edited the paper. All authors analyzed the results and reviewed the paper.

Competing interests

The authors declare no competing interests.

Additional information

Extended data is available for this paper at <https://doi.org/10.1038/s44284-025-00345-4>.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s44284-025-00345-4>.

Correspondence and requests for materials should be addressed to Linus W. Dietz.


Peer review information *Nature Cities* thanks Lincoln Larson, Catharine Ward Thompson and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.


Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025



Extended Data Fig. 1 | Pearson correlations between park health scores and the distance to the city center by city show mostly low to moderate negative correlations. This figure complements the findings regarding geographic influences on park health scores, subdivided by continents. Using a correlation

analysis between the distance to the city center (discounted using the binary logarithm) and the park health scores, we aimed to provide an additional quantification of the decay in park scores moving away from the city center.



Extended Data Fig. 2 | Inequality of park scores in the cities. Overall, nature-appreciation activities show the lowest disparities, whereas the largest differences can be observed in the social activities category. This figure visualizes the data in Table 2.

Extended Data Table 1 | Lexicon of elements and spaces for health-promoting activities in parks

Activity Category	Elements	Spaces
Cultural	information=board, tourism=artwork, historic=memorial, artwork_type=sculpture, artwork_type=statue, board_type=history, historic=monument, memorial=plaque, memorial=war_memorial, memorial=bench	tourism=artwork, tourism=attraction, religion=christian, leisure=bandstand, historic=memorial, denomination=anglican, building=church, historic=building, tourism=museum, amenity=theatre
Environmental	waste=trash, produce=plum, amenity=recycling, fruit=apple, produce=damson, leisure=garden, man_made=beehive, produce=apple, amenity=watering_place, man_made=monitoring_station	leisure=garden, landuse=flowerbed, landuse=allotments, building=greenhouse, landuse=orchard, landuse=farmland, building=farm_auxiliary, landuse=farmyard, garden:type=community, garden:type=residential
Nature-appreciation	natural=tree, amenity=fountain, tourism=viewpoint, board_type=nature, tourism=picnic_site, amenity=shelter, natural=shrub, attraction=animal, board_type=wildlife, waterway=weir	natural=wood, natural=water, natural=scrub, water=pond, natural=heath, landuse=forest, heath=bracken, natural=grassland, natural=wetland, amenity=shelter
Physical	amenity=bicycle_parking, highway=crossing, amenity=drinking_water, leisure=fitness_station, barrier=cycle_barrier, sport=fitness, leisure=pitch, sport=orienteering, orienteering=marker, leisure=playground	leisure=pitch, leisure=playground, sport=soccer, sport=tennis, highway=footway, golf=bunker, sport=basketball, highway=pedestrian, area:highway=footway, golf=tee
Social	amenity=bench, tourism=information, leisure=picnic_table, amenity=cafe, board_type=notice, amenity=telephone, amenity=fast_food, playground=playhouse, amenity=restaurant, advertising=board	amenity=cafe, building=pavilion, building=retail, amenity=community_centre, leisure=outdoor_seating, amenity=school, amenity=restaurant, building=kiosk, amenity=kindergarten, building=terrace

We show the 10 most frequent tags per activity category. The full lexicon of 1441 elements and spaces is available in the replication package.

Extended Data Table 2 | Top parks of the cities by activity category

City	Physical	Nature-appreciation	Environmental	Social	Cultural
Amsterdam	Sloterpark	Sloterpark	Park Frankendael	Westerpark	Amstelpark
Auckland	Lloyd Elmore Park	Point England Park	Paneke / Radonich Park	Ambury Regional Park	Albert Park
Berlin	Tempelhofer Feld	Landschaftspark Johannisthal/Adlershof	Gärten der Welt	Tempelhofer Feld	Treptower Park
Boston	Franklin Park	Charles River Esplanade	Temple Street Park	Charles River Esplanade	Georges Island
Buenos Aires	Parque Indoamericano	Parque de la Memoria	Paseo Arzoumanian	Parque de las Ciencias	Parque Avellaneda
Chicago	Lincoln Park	Northerly Island	Grant Park	Lincoln Park	Grant Park
Christchurch	Canterbury Agricultural Park	Bottle Lake Forest Park	Hagley Park North	Avon River Precinct	Hagley Park North
Copenhagen	Fælledparken	Østre Anlæg	Husum Bypark	Enghaveparken	Østre Anlæg
Hong Kong	九龍仔公園 Kowloon Tsai Park	藝術公園 Art Park	佐敦谷公園 Jordan Valley Park	迪欣湖活動中心 Inspiration Lake Recreation Centre	灣仔臨時海濱花園 Wan Chai Temporary Promenade
Houston	Hermann Park	Hermann Park	Wright-Bembry Park	Hermann Park	Hermann Park
London	Old Deer Park	Russia Dock Woodland	Bushy Park	Richmond Park	Alexandra Park
Madrid	Parque Agustín Rodríguez Sahagún	Dehesa de la Villa	Parque del Retiro	Finca Vista Alegre	Parque del Retiro
Melbourne	Albert Park	Grant Reserve	Fitzroy Gardens	Albert Park	Carlton Gardens
Montreal	Parc Jean-Drapeau	Parc Angrignon	Jardin botanique de Montréal	Vieux-Port	Parc Jean-Drapeau
Moscow	Парк Останкино	Бирюлёвский дендропарк	Выставка достижений народного хозяйства	Тимирязевский парк	Выставка достижений народного хозяйства
New York	Pelham Bay Park	Prospect Park	Bronx Park	Brooklyn Bridge Park	Fort Tilden
Paris	Bois de Boulogne	Bois de Vincennes	Jardin des Plantes	Bois de Boulogne	Bois de Boulogne
Perth	Altona Park	Kings Park	Hyde Park	Christ Church Grammar Playing Fields	Victoria Gardens
Philadelphia	East Fairmount Park	East Fairmount Park	East Fairmount Park	Race Street Pier	Fort Mifflin
Rio De Janeiro	Aterro do Flamengo	Quinta da Boa Vista	Largo da Carioca	Campo de Santana	Praça Luís de Camões
Rome	Villa Borghese	Villa Glori	Parco Agricolo di Casal del Marmo	Riserva Naturale dell'Acquafredda	Villa Borghese
San Diego	Balboa Park	Mission Bay Park	Balboa Park	Balboa Park	Balboa Park
San Francisco	Presidio of San Francisco	Lake Merced Park	Golden Gate Park	Presidio of San Francisco	Golden Gate Park
Seattle	Warren G. Magnuson Park	Seward Park	Washington Park Arboretum	Seattle Center	Seattle Center
Seoul	올림픽공원	매봉산공원	서울숲	송파나루공원	경복궁
Singapore	Changi Business Park	Windsor Nature Park	Singapore Botanic Gardens	The Lawn@Marina Bay	Singapore Botanic Gardens
St Petersburg	парк Героев-Пожарных	Парк-дендрарий Ботанического сада Петра Великого	Летний сад	Приморский парк Победы	парк Лесотехнической академии
Stockholm	Årstadfältet	Kungsträdgården	Sveaplan	Karlaplan	Humlegården
Sydney	Centennial Park	Centennial Park	Sydney Park	Lawrence Hargrave Reserve	Clarkes Point Reserve
Taipeh	大安森林公園	關渡自然公園	士林官邸公園	天母運動公園	中正紀念公園
Tokyo	若洲海浜公園	上野恩賜公園	北の丸公園	上野恩賜公園	上野恩賜公園
Toronto	Centennial Park	Sunnybrook Park	Highland Creek Ravine	Toronto Island Park	Don Valley Brick Works Park
Vancouver	Connaught Park	Stanley Park	Stanley Park	Hastings Park	Morton Park
Vienna	Augarten	Draschepark	Schlosspark	Schlosspark	Schlosspark
Washington DC	East Potomac Park	National Mall	Schönbrunn	Schönbrunn	Schönbrunn
			National Mall	National Mall	National Mall

We list the best-equipped park of each city by their scores on each activity category.

Extended Data Table 3 | Park health scores by quartiles determined using the distance to the city center

Activity Category	Mean Health Scores				<i>p</i> -values of H_0		
	Q1	Q2	Q3	Q4	Q1 = Q2	Q2 = Q3	Q3 = Q4
Physical	0.382	0.297	0.160	0.068	0.000	0.000	0.000
Nature-appreciation	0.107	-0.211	-0.497	-0.580	0.000	0.000	0.000
Environmental	0.020	-0.076	-0.156	-0.186	0.000	0.000	0.003
Social	0.289	0.137	-0.004	-0.088	0.000	0.000	0.000
Cultural	0.260	0.096	0.010	-0.022	0.000	0.000	0.001

Q1 are the inner city parks, and Q4 are the parks that are most distant to the city center. On the left, we show the mean health scores for each quartile; on the right, we show the *p*-values determined using a two-sided t-test for each null hypothesis $H_0: Q_i = Q_j$ stating that the mean scores for two subsequent quartiles *i* and *j* are equal. All null hypotheses can be refuted with high significance levels ($p < 0.001$, ***), with the exception of the difference between Q3 and Q4 in the environmental category, where the significance is $p = 0.003$, **. We confirmed the normal distribution of the scores using QQ-plots.

Extended Data Table 4 | The correlations between the scores for park elements and park spaces are low

Activity Category	Mean Pearson Correlation
Physical	0.16
Nature-appreciation	0.05
Environmental	0.18
Social	0.06
Cultural	0.18

This property allowed us to linearly combine them into a unified score.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a	Confirmed
☐	☒ The exact sample size (<i>n</i>) for each experimental group/condition, given as a discrete number and unit of measurement
☒	☐ A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
☐	☒ The statistical test(s) used AND whether they are one- or two-sided <i>Only common tests should be described solely by name; describe more complex techniques in the Methods section.</i>
☒	☐ A description of all covariates tested
☐	☒ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
☐	☒ A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
☐	☒ For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i>) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted <i>Give P values as exact values whenever suitable.</i>
☒	☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
☒	☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
☐	☒ Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i>), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection	The data extraction from OpenStreetmap was done using the Python bindings (https://osmcode.org/pyosmium/ v.3.6.0) for Osmium (https://osmcode.org/libosmium/)
Data analysis	The Python (v 3.9) and R (v4.5.1) code to compute park health scores is available at https://github.com/LinusDietz/Health-Promoting-Parks-Replication .

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

The replication package contains tables of the park health scores in the cities: <https://github.com/LinusDietz/Health-Promoting-Parks-Replication>. The original

OpenStreetMap data used for scoring the parks is publicly available and can be best obtained from one of the third-party download servers, for example from <https://download.geofabrik.de>. The Flickr dataset for the validation can not be shared due to terms of conditions of this dataset.

Human research participants

Policy information about [studies involving human research participants and Sex and Gender in Research](#).

Reporting on sex and gender	We have not collected any sex or gender-related information in this survey, as we did not analyze any hypotheses regarding gender, nor deemed this to be a relevant factor with respect to the goal of the survey. Keep in mind that the citizen survey is not a central part of the analysis (reported in the appendix A5 only).
Population characteristics	We collected a minimal set of demographics, namely the age groups 18-24: 16; 25-34: 35; 35-44: 15; 45-54:10; 55-64: 4; 65 and over: 1 and the postal area of the citizens. The reported postal areas were following the population distribution of London.
Recruitment	We recruited participants through the the King's College London research recruitment portal as well as mailing lists within scientific institutions in London. Thus, respondents are likely to be more educated than average citizens but encompassed varied ages as our data distribution shows. There is a potential self-selection bias, as individuals with a higher than average educational background may have been more likely to participate, which could limit the generalisability of our findings to the wider population.
Ethics oversight	King's College London Research Ethics Office, ID: MRA-22/23-38802

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☐ Behavioural & social sciences ☒ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	The study counts map objects from OpenStreetMap to quantify the overall contributions of a park towards providing means for engaging in health-promoting activities. This novel approach aims to quantify the health potential of each park in a unified score. It is an observational study design without an experimental intervention.
Research sample	The research sample were 23,477 parks in 35 global cities.
Sampling strategy	We included all named parks in all cities that were tagged as such in OpenStreetmap.
Data collection	The first author collected all data from OpenStreetMap Snapshots. These snapshots were downloaded from https://geofabrik.de , a data provider for OpenStreetMap data.
Timing and spatial scale	All data snapshots of the final analysis were downloaded on October 27, 2023. The Flickr data for the validation stems from 2004-2015.
Data exclusions	We excluded parks that did not have a name, as we expect that these parks are of insignificant size and importance.
Reproducibility	Reproducibility is possible using the provides replication pack. The software has been verified to produce the same results in repeated runs.
Randomization	Randomization is not applicable to this study design, as there are no different groups.
Blinding	As the scores are computed by a software, blinding is not applicable to this study design.
Did the study involve field work?	☐ Yes ☒ No

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology and archaeology
☒	☐ Animals and other organisms
☒	☐ Clinical data
☒	☐ Dual use research of concern

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging

Understanding the potential of urban parks to promote well-being

In the format provided by the
authors and unedited

1 Scoping Review to Map Activity Categories to Health Benefits

Our goal was to collate and map health-prompting activities in parks discussed in prior studies. Considering between a systematic and scoping type of review, the scoping review was a better fit for our task because we only needed to map activities discussed in the literature, and we did not need to focus on the types and quality of data collected in those studies, which is a task for systematic reviews. Specifically, we turned to using the well-established PRISMA method¹, which is designed to facilitate transparent reporting of reviews, and it has been designed primarily for reviews of studies that evaluate the effects of health interventions, irrespective of the design and strength of effects found in the included studies.



Suppl. Fig. 1. Our PRISMA Statement: Process of identification, screening, and determining eligibility for articles in our literature survey.

The overarching research question was: “Which are the health benefits of activities in urban greenery?” Our focus on urban greenery instead of only parks was to ensure both the *comprehensiveness* and *generality* of the taxonomy, as future studies might look beyond urban parks. We used the WHO’s definition of urban greenery to determine the scope of our survey: “[...] urban green space is defined as all urban land covered by vegetation of any kind. This covers vegetation on private and public grounds, irrespective of size and function, and can also include small water bodies such as ponds, lakes, or streams (“blue spaces”)².”

As we were interested in the intersection of urban greenery and medical studies, we performed a set of queries on PubMed and SpringerLink to identify papers that linked the usage of urban greenery with health benefits. An article was deemed relevant if the results evidenced that one or more activities typically done in public urban green spaces had a health benefit. To obtain a comprehensive overview of each activity category, we used a total of 6 queries. Upon our preliminary experiments, we employed a collection of keywords for our queries that included both those commonly encountered in the initial set of studies and those formulated by our experts. This approach enabled us to discover a diverse range of papers relevant to each category of activity. The queries were:

Physical activities: (urban greenery) AND (health) AND (sports OR exercise)

Nature-appreciation activities: (urban greenery) AND (health) AND (nature) AND (exposure)

Environmental activities: (urban greenery) AND (health) AND (garden OR planting OR conservation)

Social activities: (urban greenery) AND (health) AND (social OR social cohesion OR social capital OR social contacts)

Cultural activities: (urban greenery) AND (health) AND (culture) OR (cultural ecosystem)

Mind-body activities: (urban greenery) AND (health) AND (mindfulness OR meditation OR yoga OR tai chi OR breathing techniques)

Following the PRISMA statement depicted in [Suppl. Fig. 1](#), out of the initially *identified* 762 articles, 417 were duplicates, leaving us with 344 unique articles. Next, we *screened* these articles and discarded 5 conference proceedings and 17 articles that were perspectives or study protocols, successfully retrieving the remaining 322 articles. In the *Eligibility* step, we determined

Suppl. Tab. 1. Activities in urban parks linked to health benefits. Specific health benefits evidenced in the respective articles are grouped by health aspects.

Activity Category	Health Aspect	Specific Health Benefit
Physical	Cognitive health	dementia prevention ³
	General health	longevity ^{4,5}
	Mental health	stress reduction ^{6–10} , depression prevention ^{10–17} , anxiety reduction ¹⁴ , various ¹⁸ , mood improvement ¹⁹
	Physical health	weight reduction ^{4,20–26} , increase of physical activity ^{7,9,13,15,27–49} , blood pressure reduction ¹⁶ , diabetes prevention ^{4,50} , various ¹⁸ , increase of leisure activities ⁵¹ , hypertension ⁴ , cardiovascular health improvements ^{24,50,52–54} , bone development ²⁴
	Social health	various ^{18,55} , social cohesion ^{13,16,56,57}
	Well-being	increase restorative capacity ⁵⁸ , enhanced social interactions ⁵⁹ , quality of life ^{3,15,37,44,54}
Nature-appreciation	Cognitive health	attention fatigue reduction ⁶⁰
	General health	lower morbidity ⁶¹
	Mental health	positive emotions ^{62,63} , depression prevention ^{12,13,15,16,64–67} , anxiety reduction ^{65,66,68,69} , suicide prevention ^{70,71} , mood improvement ^{19,60,67,72} , relaxation ⁶³ , mindfulness ⁶³ , calmness ⁶⁹ , stress reduction ^{7,60,65,66,72–78}
	Physical health	mood improvement ⁷² , improved ghq-12 scores ⁷⁹ , blood pressure reduction ^{16,72,80} , antenatal health ⁸¹ , respiratory health ⁸² , increase of physical activity ^{7,13,15,46,79,83,84} , blood oxygen saturation ⁷²
	Social health	increased social capital ⁸⁵ , social loneliness reduction ⁸⁶ , various ⁵⁵ , social cohesion ^{13,16}
	Well-being	stress reduction ^{65,72,77} , quality of life ^{15,64,65,77,87–89} , blood pressure reduction ⁷² , increase restorative capacity ⁹⁰
Environmental	Cognitive health	restorative effect against cognitive failures ⁹¹
	General health	lower morbidity ⁶¹
	Mental health	stress reduction ^{76,92} , anxiety reduction ^{93,94} , improved sleep ^{93,94} , depression prevention ^{64,94}
	Physical health	cardiovascular health improvements ⁹⁵ , inflammation reduction ⁹¹ , respiratory health ⁹⁵ , access to healthy produce ⁹⁶ , immune system improvement ⁹⁷ , increase of physical activity ^{29,98} , improved sleep ⁹³
	Social health	access to healthy produce ⁹⁶ , social cohesion ^{92,96}
	Well-being	nutritional diversity ⁹⁹ , quality of life ^{64,93,100,101} , improved sleep ⁹³ , increase restorative capacity ¹⁰²
Social	Cognitive health	dementia prevention ³ , restorative effect against cognitive failures ⁹¹
	General health	longevity ⁵
	Mental health	mood improvement ¹⁰³ , depression prevention ^{10,11} , various ¹⁰⁴ , improved mental health inventory (mhi-5) scores ¹⁰⁵ , stress reduction ^{10,92}
	Physical health	access to healthy produce ⁹⁶ , various ¹⁰⁴ , inflammation reduction ⁹¹ , increase of physical activity ^{44,49}
	Social health	social cohesion ^{56,57,92,96,103,106} , various ⁵⁵ , access to healthy produce ⁹⁶ , increased social capital ¹⁰⁷ , social loneliness reduction ¹⁰⁸ , improve sense of social belonging ^{109–111}
	Well-being	increase restorative capacity ¹⁰² , quality of life ^{3,44,112} , enhanced social interactions ⁵⁹
Cultural	Cognitive health	dementia prevention ³
	General health	various ¹¹³
	Physical health	increase of physical activity ¹¹⁴
	Well-being	quality of life ^{3,112,114}
Mindfulness	Mental health	stress reduction ⁷⁴ , anxiety reduction ¹⁴ , depression prevention ^{14,17}
	Physical health	increase of physical activity ⁴⁹
	Well-being	quality of life ¹¹⁵

whether these articles were relevant to our search. We found that 114 articles were relevant, while 208 were not. Most articles were excluded because they were not about urban green spaces or because there was no significant link between the activities and health benefits. When analyzing the *included* articles, we recorded each activity category alongside the general health aspects and specific health benefits the article evidenced (Suppl. Tab. 1).

Results

As can be seen in Suppl. Tab. 1, most research has focused on the benefits of *physical activity* in parks. Out of 79 studies on the health benefits of exercising, 46 underscored positive outcomes like weight loss^{20–22}, cardiovascular improvements^{50,52}, metabolic activity^{28,29}. Additionally, these activities demonstrated positive effects on mental health (16 articles), well-being (7 articles), and social health (6 articles).

The second most studied category is *nature-appreciation*, with 68 articles. These activities significantly boost mental health (34 articles), primarily in reducing stress^{7,73} and anxiety^{65,68} and preventing depression^{12,13,15,16,65–67}. They also contribute to physical health (14 articles) and overall well-being (12 articles).

In our review, we found that *social* and *environmental activities* received less attention in conjunction with urban parks, with only 33 and 28 articles covering them, respectively. Despite this, both contribute to all identified health aspects. Social activities enhance social and mental health, fostering a sense of belonging^{109–111} and improving mood¹⁰³. Environmental activities, such

as gardening, offer diverse benefits, including cognitive restoration⁹¹ and improved general health⁶¹.

Finally, *cultural* and *mindfulness activities* are relatively under-researched regarding their health benefits in the context of urban parks. Cultural activities often fell outside the scope of our review, which required a connection to urban parks, while more general cultural activities were studied. However, their health benefits are likely underreported given the presence of cultural facilities in parks (e.g., historic monuments or arts venues). Likewise, mindfulness activities and nature-based therapeutic interventions, such as forest therapy¹¹⁶ provide health benefits, but have not been studied in the context of urban parks, highlighting a potential gap in the scientific literature that warrants future exploration.

In summary, we found that all the different activities we identified have a *distinct but overlapping set of health benefits*. For example, both physical and nature-appreciation activities help prevent depression, but only physical activities help bone development, and only nature-appreciation brings calmness.

2 Validating the Overall Ranking of Parks Through an Online Survey

As an additional means for validating the park scores, we conducted a survey in one city, London, UK. In an online questionnaire, we asked London citizens about suitable parks for performing activities. The main set of questions was phrased as: “*Can you name several parks suitable for physical activities (e.g., sports)?*”

Study Information

We recruited the participants using the first author’s institutional research recruitment portal as well as mailing lists within scientific institutions in London. The participants were informed about the voluntary nature of their participation and that no personally identifiable information about them was collected. For these reasons, age was collected using 7 age groups (“*Below 18*”, “*18–24*”, “*25–34*”, “*35–44*”, “*45–54*”, “*55–64*”, “*65 and over*”) and as a privacy mechanism only the postal area (e.g., N1) instead of the full postcode was requested. Furthermore, we asked participants how long they have been living in London (“*I don’t live in London.*” – “*Less than 1 year.*” – “*1 to 5 years.*” – “*More than 5 years.*”). Finally, as a means to identify low-quality responses, we asked people for a park close to their homes, which we could use as an instructional manipulation check in conjunction with the reported postal area. The data collection was registered as a minimal-risk study at the first author’s institutional review board (King’s College London Research Ethics Office, ID: MRA–22/23–38802).

Results


The metric we used to quantify how well the citizen response aligned with our health scores is the average percentile-ranking^{117,118}, which captures how highly the selected park was placed in the overall ranking of parks for the corresponding activity. A value close to 1 means parks with the highest scores were selected, whereas 0.5 would represent a random selection. The results demonstrate a clear alignment between the freely recalled parks by the participants and the rankings derived from our health scores. As shown in [Suppl. Tab. 2](#), the median and mean values of the average percentile-ranking for the parks named by citizens were consistently high. For nature-appreciation, physical activities, cultural activities, and social activities the median scores are above 0.89, highlighting a strong concordance between citizens’ perceptions of the park and the quantitative rankings derived from our proposed park profiling method.

Suppl. Tab. 2. Result statistics of the online survey. Citizens were asked to name parks that are suitable for the activities. The first three columns show the statistics of the average percentile-ranking of the named parks. **AR** is the answer rate of the respective category, i.e., how many respondents were able to name at least one park), **N** is the number of non-empty responses, and **MR** is the mean number of parks that were named per respondent.

Activity category	median	mean	σ	AR	N	MR
Physical	0.91	0.84	0.17	97.5%	78	4.26
Nature-appreciation	0.95	0.85	0.19	95%	76	2.75
Environmental	0.50	0.50	0.31	57.5%	46	1.80
Social	0.93	0.87	0.13	96.2%	77	4.94
Cultural	0.89	0.75	0.25	81.2%	65	2.27

The result for environmental activities is subpar compared to the other activities, with a mean and median average percentile-ranking of 0.50. Only 57.5% of the respondents could name an environmental park, and on average, 1.8 parks were named in this category by each person, which indicates that parks for environmental activities are harder to think of compared to the other activities. Another explanation for the low scores in this category is that while urban gardening and conservation can be done in many parks, they typically do not occupy much *spaces*, environmental activities are less mainstream in cities, and the number of *park elements* for this category is comparatively low in London’s parks impeding high activities scores.

The overall alignment between Londoners' perceptions of parks and our health scores underscores the effectiveness of our approach in accurately capturing and evaluating the health-promoting potential of parks.




Suppl. Fig. 2. The online survey results as box-whisker plots. Standard box plot with Q1, median, Q3; whiskers at $1.5 \times$ interquartile range. The values indicated with the x represent the mean values.

3 Supplementary Methodological Details

3.1 Determining the Threshold Values for Computing the Linear Models

Suppl. Fig. 3 and Suppl. Fig. 4 depict histograms of park elements and park spaces. The plots supplement the determination of thresholds for excluding parks with insufficient activity data.




Suppl. Fig. 3. Histogram of park elements. We set 2 as the minimum number per activity category.

3.2 Data Cleaning of OSM Tags

This section describes the steps we undertook to exclude OSM tags that are not useful for our analysis.

The first step of the data cleaning process was not specific to annotating health-promoting activities. Instead, the focus was on removing any extra information not necessary for understanding the main purpose of the map object. For example, to identify a bench on a map, one just needs to look for the label `amenity=bench`. However, a bench can also have additional tags like `inscription`, `operator`, `material`, and `backrest`, which offer more specifics about the bench. When it comes to identifying the object's primary purpose for health-promoting activities, this extra information is not only unnecessary but could also lead to confusion. To remove these irrelevant labels, three of the authors created lists of keys and values that



Suppl. Fig. 4. Histogram of park spaces. We set 0.05ha (500 m²) as the minimum size per activity category.

were only used to provide extra details when combined with other labels. All co-authors carefully reviewed, discussed, and agreed upon these lists. If there was any doubt about whether to exclude certain labels, they were kept and left for subsequent annotation. The goal was to make sure that only necessary and relevant labels were kept for categorizing *park elements* and *spaces* into health-promoting activity categories.

In the process of cleaning the data, 1926 keys were omitted. These included keys such as `name`, `operator`, and `source`, which cannot provide insight into the object's activity. In addition, 11 values were also left out because they only described metadata and did not help in understanding the primary function or essence of the map object. Examples of such values include `yes/no`, `unknown`, or `Bing`. A full list of these omitted keys and values can be found in the replication repository. This initial data cleaning step significantly reduced the number of tags to 2118, which were the ones we needed to map to health-promoting activities, or none if the object did not support any of them. This streamlined dataset provided a more focused and relevant basis for the subsequent annotation and classification of *park elements* and *spaces*.

3.3 Benchmarking LLM Classifiers

To evaluate the suitability of large language model (LLM) classifiers as annotators for OpenStreetMap (OSM) tags, we created a high-quality, expert-annotated gold standard set consisting of the 100 most frequent tags. To ensure accuracy and reliability, three co-authors independently labeled these 100 items with health-promoting activities or none, and we used the majority voting strategy to aggregate the individual opinions into one final outcome label. In cases where conflicts arose, i.e., where the three annotators provided different labels, a discussion was held to resolve the discrepancies. Through this rigorous annotation process, we established a robust and reliable “*gold standard*” dataset of 100 items. This dataset serves as a benchmark to assess the accuracy of the labels provided by the LLM classifiers.

For generating the annotations, we conducted a systematic exploration of the configuration settings of two LLMs, GPT-3.5-turbo¹¹⁹ and GPT-4¹²⁰ using the OpenAI API¹²¹. Our goal was to identify the best-performing setting in terms of the weighted F_1 score, which is the harmonic mean of precision and recall in this multi-class classification task. The independent variables were *i*) the large language model, i.e., `gpt-3.5-turbo` or `gpt-4`¹²⁰, *ii*) the temperature parameter $t \in \{0.3; 0.6; 0.9\}$, which controls the randomness of the models' completions, and *iii*) the prompt, for which we tested two versions, one with and without providing a brief definition of the OSM tag taken from the OSM wiki. The full prompt is shown in Suppl. Fig. 5.

Suppl. Fig. 5 shows a specific sequence of prompts designed to elicit a main activity and a secondary activity for each OSM tag. The reason behind this approach was our hypothesis that certain OSM tags could support multiple health-promoting activities, as demonstrated by the example of benches that could be argued to be annotated with social, nature-appreciation, or physical activities. Additionally, we obtained a reliability score for each of the model's annotations. These reliability scores offer an indication of the model's confidence in its assigned activities, which could serve as a threshold to actually use the annotations, as low scores might indicate that the annotation is more speculative. By incorporating these main activities, secondary activities, and reliability scores from the LLM models, we hoped to gain a more nuanced insight into how these amenities and facilities in parks can be used. This detailed information allowed us to account for the potential multi-functionality of certain OSM tags and provided data for the evaluation using the proposed benchmark.

Furthermore, we followed the guidelines¹²² to optimize the performance of the LLMs annotations. We assigned a system persona, i.e., ‘*You are an expert in urban planning and public health, with a specialization in urban parks. [...]*’, gave definitions of the six activities with exemplary activities, and provided several correct completions of items as means to few-shot learning. Finally, we provided a clear specification of the desired output format.

To determine the highest agreement between the human-annotated benchmark and the annotations of the LLMs, we used the

Suppl. Tab. 3. Results of LLM benchmarking. The highest performance is achieved with GPT-4, using a temperature of 0.9 and not providing definitions for the tags. Annotating a secondary activity did not improve the F_1 scores.

LLM	Definitions	Temperature	F_1 -score Main Activity	F'_1 -score Weighted Combination
gpt-4	✗	0.9	0.772	0.772
gpt-4	✗	0.6	0.770	0.770
gpt-4	✓	0.3	0.764	0.764
gpt-4	✗	0.3	0.755	0.755
gpt-3.5-turbo	✗	0.6	0.747	0.747
gpt-4	✓	0.6	0.740	0.740
gpt-3.5-turbo	✗	0.9	0.728	0.728
gpt-3.5-turbo	✓	0.3	0.726	0.726
gpt-4	✓	0.9	0.713	0.713
gpt-3.5-turbo	✓	0.6	0.710	0.710
gpt-3.5-turbo	✗	0.3	0.704	0.708
gpt-3.5-turbo	✓	0.9	0.689	0.689

F_1 score, which is the harmonic mean of the precision and recall. One complication in the evaluation was that the benchmark only comprised one activity label, whereas we asked the LLM annotator for a main and secondary activity for each tag. Thus, we report two F_1 scores: one that uses the label from the main category only and another that is a weighted combination of the main activity category and the secondary activity category. The weighted F'_1 score is computed by slightly altering the impact of each element of the confusion matrix as follows:

$$TP' = TP_{main}^i \cdot reliability^i + TP_{2nd}^i \cdot (1 - reliability^i) \quad (\text{true positives}) \quad (1)$$

$$FP' = FP_{main}^i \cdot reliability^i + FP_{2nd}^i \cdot (1 - reliability^i) \quad (\text{false positives}) \quad (2)$$

$$FN' = FN_{main}^i \cdot reliability^i + FN_{2nd}^i \cdot (1 - reliability^i) \quad (\text{false negatives}) \quad (3)$$

$$reliability^i = \frac{mean(reliability_{main}^i)}{mean(reliability_{main}^i) + mean(reliability_{2nd}^i)} \quad (\text{Ratio of reliability between main and secondary category}) \quad (4)$$

Intuitively, this means that we use the reliability scores stemming from the LLM annotations to estimate the LLM's confidence that a label is correct, thus creating a comparable metric that allows for comparing two annotations for one item to one human-annotated ground truth.

We tested various settings to see which would deliver the best performance, which was GPT-4, set at a temperature of 0.9, and without providing definitions for tags. To give you a clearer picture, we've compiled the results of the top-performing configuration in [Suppl. Tab. 3](#). The F_1 scores tabulated in the tables show the best results of systematically adjusting the reliability scores for primary and secondary categories from 0 (using any label, regardless of its reliability) to 1 (annotate "none" in all cases). Generally, GPT-4 outperformed its predecessor, GPT-3.5. Adding definitions actually had a negative effect on label quality, possibly due to misleading keywords in the tagging instructions. When it came to the temperature setting, there was no consistent impact, with minimal differences between otherwise equivalent configurations. Interestingly, adding a secondary activity label didn't improve the annotation quality (cf. last column of [Suppl. Tab. 3](#)). In fact, the best results were achieved when the reliability threshold of the secondary annotation was close to 1, rendering all secondary annotations to "none", thus being equivalent to only using the main activity label. This suggests that the primary labels generated by the system are already of high quality, so putting any weight on a secondary label actually harms the overall score. Based on these findings, we decided to use GPT-4, set at a temperature of 0.9 and without definitions, to label all OSM tags and not impose any threshold on the reliability score.

3.4 Modeling Average Park Offerings

To model average park offerings, we used independent regression models for *park elements* and *spaces*. During the computation of the regression models, parks with very low activity counts in a specific category were excluded. This exclusion was necessary to prevent artificially flattening the regression lines due to close-to-zero values, which would distort the normalization. The specific threshold for excluding parks with low activity counts was determined empirically by analyzing the histograms of the values. This approach enabled us to identify an appropriate cutoff point for excluding parks with insufficient activity data, ensuring the reliability of the regression models. For a visual representation of the exclusion process and the determination of the threshold, refer to [Suppl. Fig. 3](#) and [Suppl. Fig. 4](#).

3.5 Orthogonality of Park Elements and Spaces

Park elements include points of interest, individual trees, benches, and similar items. *Park spaces*, however, include areas like forests, sports fields, and buildings. There can be cases where a park area is broken down into its individual parts, like a playground with separately mapped features like swings, slides, or spinning equipment. But these cases are pretty rare in OSM mapping. Likewise, unless a tree is particularly important, areas tagged as `natural=wood` should not include individual trees according to the mapping guidelines¹²³. Based on these observations, we hypothesized that it would be acceptable to combine scores from park features and areas linearly, as they contribute differently to the overall offering of facilities for health-promoting activities.

To validate the assumption that *park spaces* and *park elements* are orthogonal, we calculated the pairwise correlation coefficients of their respective scores in all cities and averaged them, as presented in Extended Data Table 4. The low Pearson Correlation Coefficients supported our observation that the scores of *park spaces* and *park elements* indeed capture largely independent concepts, with all correlations being slightly positive but below 0.2. Consequently, we combined them into one overall score for the park.

3.6 Semantic Matching of Flickr Labels and OSM Tags

In our validation, we used a global dataset of geotagged photos from Flickr, from which we selected all 10,711,513 images that were taken within one of the parks from 35 cities in our study. These images came with user-generated labels and were also partially annotated with computer vision labels from a computer vision algorithm^{124,125}. To obtain semantically equivalent representations of Flickr labels and OSM tags, we employed Sentence-BERT (S-BERT¹²⁶) for text embeddings. We formulated this task as an asymmetric semantic search problem, where the Flickr label was the search term, and the goal was to find the closest matching OSM tag. Given the worldwide reach of our study, the multiple languages present in the user-generated Flickr labels created a challenge in mapping them to the corresponding OSM tags, which were all in English. To address this, we identified the top three languages besides English used in the tags of each city, using the Google MediaPipe¹²⁷ Language Detection Model¹²⁸. To ensure that the language detection was accurate and to eliminate named entities, we only used labels where the language detection indicated a confidence of 50% or more. We then translated those tags to English using the respective OPUS machine translation models¹²⁹.

To further improve the quality of embeddings, we augmented the OSM tags with short definitions sourced from the OSM mapping guidelines¹²³. For instance, the OSM tag `sport=table_tennis` was augmented with the definition "A bat and ball game played over a table." We were able to expand 66% of the OSM tags with these descriptions. The remaining tags were left without descriptions primarily because of the unregulated nature of tagging in OSM, which led to many undocumented tags or multiple values within one tag, like `sports=soccer;rugby`. Note that these tags were still used for mapping, albeit with less information.

After embedding the OSM tags using S-BERT's `all-mpnet-base-v2` model, we proceeded to match each Flickr label to the closest OSM tag in the embedding space, using the cosine distance as similarity measure. To ensure that the matches were of high quality, we set a strict threshold: the cosine similarity score had to be at least 0.7. We arrived at this value after noticing that when the similarity score was lower than 0.7, the matches became less reliable based on manual inspections. This allowed us to avoid matching labels that did not have meaningful OSM counterparts. For example, abstract labels describing certain phenomena like "cloud", "rain", and "sunset" were not matched.

A detailed review of the matched pairs revealed that, as anticipated, most pairings were logical based on the text similarity between labels and tags with definitions. However, some minor adjustments were still needed, as some matches were not entirely consistent with the theme of health-promoting activities in parks. For example, the term "outdoor" was initially linked to `swimming_pool=outdoor`. But as there cannot be a suitable equivalent for "outdoor" on OSM, we removed this pairing and equivalent ones, such as "park," as all photos were taken in parks. Another instance was the pairing of "water", which did not capture the specific role of water features in parks in promoting health. We manually adjusted this to `water=river`, which better reflects bodies of water commonly found in parks. Through this review step, we improved the quality of the matched pairs, ensuring they more closely align with the theme of health-promoting park activities. The need for this manual step should not diminish the effectiveness of the semantic search within sentence embeddings. It was merely to eliminate labels

that could not meaningfully correspond with an OSM tag and to match a few labels with more domain-relevant tags. This matching process yielded 2,171 label-tag pairs in total. Of these, 1,432 pairs corresponded to an OSM tag with health-promoting benefits, such as “*steeplechase*” being matched to `athletics=steeplechase` involving physical health benefits, while 739 pairs, such as “*Lamp Post*” being matched to `man_made=lamp_post` did not imply health benefits.

We evaluated the accuracy of the resulting label-tag matchings by asking three domain experts to independently assess whether the 20 most frequent matchings from Flickr tags to activity categories were plausible and correct. We aggregated their responses using majority voting. Given the multiple languages present in the dataset, we used only the tags from London in this evaluation step, as they were in English. The experts agreed with 82% of the matchings, which is highly accurate considering they are based solely on individual tags.

Having assured that the matchings are accurate, we proceeded to profile the parks based on the activities associated with the matched OSM tags, following the same scoring approach as what we used for the OSM *park elements* and *spaces*. In our validation, we chose a minimum of 250 images from each park and at least 15 parks in each city. This criterion was established to secure a robust number of images for each park, enhancing the accuracy of our analysis. This approach helped us avoid any potential bias that could have been introduced by individual photographers if a park had only a few images.

Suppl. Fig. 6 depicts the computation of the Flickr activity scores. The method is the same as for the OSM tags; however, on the y-axis, we use the count of the matched Flickr labels instead of the OSM tags.

3.7 User Studies with Urban Designers and Park Maintenance Experts

We conducted three semi-structured interviews, each lasting about 30 minutes, with urban designers and park maintenance experts. Participants were recruited through our network and by direct email to municipalities. The experts’ backgrounds were as follows:

- Lecturer and urban designer (E1). 20 years of practical and academic experience in Hong Kong and London.
- Urban designer and master planner (E2). 30 years of experience in the United States, United Kingdom, Europe, and the Middle East.
- Parks development manager at a local council (E3). 12 years of experience as conservationalist, arboricultural manager in diverse contexts.

The online interviews consisted of 4 steps.

1. **Introduction.** Three questions to understand their experience in urban park design and management.
2. **Study brief.** A summary of our study and main findings.
3. **Visualization.** Interacting with the visualization and exploring a chosen city.
4. **Debriefing.** Open-ended questions and feedback on the project.

We qualitatively analyzed the interview transcripts^{130,131} (recorded with consent) to understand the implications of our work for professional practice, verify inter-cultural validity, and to further validate the taxonomy of activities.

⇒ You are an expert in urban planning and public health, with a specialization in urban parks. You have studied how parks promote health and have an understanding of the various activities that people engage in within them. Proficient in the OpenStreetMap project and skilled in tagging urban elements, particularly those within parks, your responsibility involves assigning activities to specific park elements based on OpenStreetMap tags.

⇒ Consider these 6 categories of activities people do in urban parks:

Physical activities This category is about leisure pursuits that involve physical movement and sports. Example activities are: Walking, hiking, trail running, biking, swimming, rock climbing, canoeing, kayaking, horseback riding, outdoor sports, and group fitness classes.

Mind-body activities This category is about physical practices that combine movement, breathing techniques, and meditation to promote relaxation, stress reduction, and overall well-being. Example activities are: Yoga, meditation, and tai chi.

Nature appreciation activities This category is about leisure pursuits that involve enjoying and exploring the natural world. Example activities are: Bird watching, camping, picnicking, fishing, painting, drawing, photography, and nature journaling.

Environmental activities This category is about gardening and conservation of parks. Example activities are: Gardening, planting trees and flowers, and participating in conservation efforts and volunteering.

Social activities This category is about coming together and communal experiences that involve engaging with others. Example activities are: Attending outdoor festivals and concerts, playing sports, and volunteering.

Cultural activities This category is about activities that celebrate and showcase the cultural diversity and heritage of the community. Example activities may include cultural festivals, art exhibits, music performances, storytelling, and cultural workshops.

Given the categories of "Physical activities", "Mind-body activities", "Nature appreciation activities", "Environmental activities", "Social activities", "Cultural activities", and "none of the above"; How would you classify the following OpenStreetMap Tags found in a park and which [0,100] reliability scores (even multiple scores) would you give to your assessment? You can assign up to two categories, but you don't have to!

⇒ Provide the answer in the format: Main Activity|Reliability Score (of the main activity)|Secondary Activity (if it exists)|Reliability Score (of the secondary activity)|Explanation (not more than 20 words)

⇒ leisure=playground

⇒ Social|80|Physical|70|Children utilize playgrounds for both physical activity and exercise, as well as for social interactions through cooperative play.

⇒ leisure=pitch

⇒ Physical|90|Social|75|People engage in team sports on sports pitches, encompassing both physical activity and a social aspect due to the collaborative nature of team-based sports.

⇒ leisure=garden

⇒ Nature-appreciation|80|Environmental|60|People utilize gardens to cultivate an appreciation for nature, while gardens also serve as spaces for environmental activities, such as planting flowers and engaging in gardening practices.

⇒ amenity=bench

⇒ Social|80|Nature-appreciation|40|People utilize benches for relaxation and social interaction, fostering a sense of community. Moreover, benches offer the opportunity to immerse oneself in the surrounding nature, encouraging moments of contemplation and introspection.


⇒ natural=wood

⇒ Nature-appreciation|100|None|100|Forested areas, such as woods, primarily serve the purpose of appreciating and connecting with nature.

⇒ amenity=parking_space

⇒ None|90|None|100|A parking space does not inherently cater to a specific activity.

Suppl. Fig. 5. Preparatory prompt provided to the LLM classifiers via the OpenAI API. The tag and the definition were subsequently prompted. Regular text refers to 'user' messages, gray text refers to 'system' messages, and underlined text refers to 'assistant' messages. ⇒ denotes the beginning of a new message. Bold markup was added for improved readability.



Suppl. Fig. 6. Visualizing the fit of the linear model for determining the park scores using matched Flickr labels for London, UK. The horizontal axis denotes the park's area ($\log_2$), and the vertical axis is the number of categorized labels of images from these parks ($\log_2$).

References

1. Page, M. J. *et al.* The prisma 2020 statement: an updated guideline for reporting systematic reviews. *BMJ: Br. Med. J.* n71, DOI: [10.1136/bmj.n71](https://doi.org/10.1136/bmj.n71) (2021).
2. World Health Organization. Regional Office for Europe. *Urban green spaces: a brief for action* (World Health Organization. Regional Office for Europe, Copenhagen, Denmark, 2017).
3. Röhr, S. *et al.* How can urban environments support dementia risk reduction? a qualitative study. *Int. J. Geriatr. Psychiatry* **37**, DOI: [10.1002/gps.5626](https://doi.org/10.1002/gps.5626) (2021).
4. Aliyas, Z. Built environment correlates of walking for recreation or exercise. *J. Public Heal.* **27**, 349–356, DOI: [10.1007/s10389-018-0956-y](https://doi.org/10.1007/s10389-018-0956-y) (2018).
5. Takano, T., Nakamura, K. & Watanabe, M. Urban residential environments and senior citizens' longevity in megacity areas: the importance of walkable green spaces. *J. Epidemiol. & Community Heal.* **56**, 913–918, DOI: [10.1136/jech.56.12.913](https://doi.org/10.1136/jech.56.12.913) (2002).
6. Watson, C. *et al.* The association between natural outdoor environments and common somatic symptoms. *Heal. & Place* **64**, 102381, DOI: [10.1016/j.healthplace.2020.102381](https://doi.org/10.1016/j.healthplace.2020.102381) (2020).
7. Ribeiro, A. I. *et al.* Exposure to nature and mental health outcomes during COVID-19 lockdown. a comparison between portugal and spain. *Environ. Int.* **154**, 106664, DOI: [10.1016/j.envint.2021.106664](https://doi.org/10.1016/j.envint.2021.106664) (2021).
8. Lin, W. *et al.* Sitting or walking? analyzing the neural emotional indicators of urban green space behavior with mobile EEG. *J. Urban Heal.* **97**, 191–203, DOI: [10.1007/s11524-019-00407-8](https://doi.org/10.1007/s11524-019-00407-8) (2020).
9. de Vries, S., van Dillen, S. M., Groenewegen, P. P. & Spreeuwenberg, P. Streetscape greenery and health: Stress, social cohesion and physical activity as mediators. *Soc. Sci. & Medicine* **94**, 26–33, DOI: [10.1016/j.socscimed.2013.06.030](https://doi.org/10.1016/j.socscimed.2013.06.030) (2013).
10. Chen, K., Zhang, T., Liu, F., Zhang, Y. & Song, Y. How does urban green space impact residents' mental health: A literature review of mediators. *Int. J. Environ. Res. Public Heal.* **18**, 11746, DOI: [10.3390/ijerph182211746](https://doi.org/10.3390/ijerph182211746) (2021).
11. Wang, R., Liu, Y., Xue, D. & Helbich, M. Depressive symptoms among chinese residents: how are the natural, built, and social environments correlated? *BMC Public Heal.* **19**, DOI: [10.1186/s12889-019-7171-9](https://doi.org/10.1186/s12889-019-7171-9) (2019).
12. Annerstedt, M. *et al.* Green qualities in the neighbourhood and mental health – results from a longitudinal cohort study in southern sweden. *BMC Public Heal.* **12**, DOI: [10.1186/1471-2458-12-337](https://doi.org/10.1186/1471-2458-12-337) (2012).
13. Cox, D. *et al.* Doses of nearby nature simultaneously associated with multiple health benefits. *Int. J. Environ. Res. Public Heal.* **14**, 172, DOI: [10.3390/ijerph14020172](https://doi.org/10.3390/ijerph14020172) (2017).
14. Bray, I., Reece, R., Sinnett, D., Martin, F. & Hayward, R. Exploring the role of exposure to green and blue spaces in preventing anxiety and depression among young people aged 14–24 years living in urban settings: A systematic review and conceptual framework. *Environ. Res.* **214**, 114081, DOI: [10.1016/j.envres.2022.114081](https://doi.org/10.1016/j.envres.2022.114081) (2022).
15. Gianfredi, V. *et al.* Association between urban greenspace and health: A systematic review of literature. *Int. J. Environ. Res. Public Heal.* **18**, 5137, DOI: [10.3390/ijerph18105137](https://doi.org/10.3390/ijerph18105137) (2021).
16. Shanahan, D. F. *et al.* Health benefits from nature experiences depend on dose. *Sci. Reports* **6**, DOI: [10.1038/srep28551](https://doi.org/10.1038/srep28551) (2016).
17. Marx, W. *et al.* Clinical guidelines for the use of lifestyle-based mental health care in major depressive disorder: World federation of societies for biological psychiatry (WFSBP) and australasian society of lifestyle medicine (ASLM) taskforce. *The World J. Biol. Psychiatry* **24**, 333–386, DOI: [10.1080/15622975.2022.2112074](https://doi.org/10.1080/15622975.2022.2112074) (2022).
18. Sallis, J. F. *et al.* Co-benefits of designing communities for active living: an exploration of literature. *Int. J. Behav. Nutr. Phys. Activity* **12**, DOI: [10.1186/s12966-015-0188-2](https://doi.org/10.1186/s12966-015-0188-2) (2015).
19. Aspinall, P., Mavros, P., Coyne, R. & Roe, J. The urban brain: analysing outdoor physical activity with mobile EEG. *Br. J. Sports Medicine* **49**, 272–276, DOI: [10.1136/bjsports-2012-091877](https://doi.org/10.1136/bjsports-2012-091877) (2013).
20. Li, X. & Ghosh, D. Associations between body mass index and urban “green” streetscape in cleveland, ohio, USA. *Int. J. Environ. Res. Public Heal.* **15**, 2186, DOI: [10.3390/ijerph15102186](https://doi.org/10.3390/ijerph15102186) (2018).
21. Zang, P. *et al.* Effect of built environment on BMI of older adults in regions of different socio-economic statuses. *Front. Public Heal.* **11**, DOI: [10.3389/fpubh.2023.1207975](https://doi.org/10.3389/fpubh.2023.1207975) (2023).
22. Kim, J.-H., Lee, C., Olvera, N. E. & Ellis, C. D. The role of landscape spatial patterns on obesity in hispanic children residing in inner-city neighborhoods. *J. Phys. Activity Heal.* **11**, 1449–1457, DOI: [10.1123/jpah.2012-0503](https://doi.org/10.1123/jpah.2012-0503) (2014).

23. Bourdeaudhuij, I. D. *et al.* International study of perceived neighbourhood environmental attributes and body mass index: IPEN adult study in 12 countries. *Int. J. Behav. Nutr. Phys. Activity* **12**, DOI: [10.1186/s12966-015-0228-y](https://doi.org/10.1186/s12966-015-0228-y) (2015).
24. Veitch, J. *et al.* Understanding children's preference for park features that encourage physical activity: an adaptive choice based conjoint analysis. *Int. J. Behav. Nutr. Phys. Activity* **18**, DOI: [10.1186/s12966-021-01203-x](https://doi.org/10.1186/s12966-021-01203-x) (2021).
25. Mackenbach, J. D. *et al.* Obesogenic environments: a systematic review of the association between the physical environment and adult weight status, the SPOTLIGHT project. *BMC Public Heal.* **14**, DOI: [10.1186/1471-2458-14-233](https://doi.org/10.1186/1471-2458-14-233) (2014).
26. Daniels, K. M., Schinasi, L. H., Auchincloss, A. H., Forrest, C. B. & Roux, A. V. D. The built and social neighborhood environment and child obesity: A systematic review of longitudinal studies. *Prev. Medicine* **153**, 106790, DOI: [10.1016/j.ypmed.2021.106790](https://doi.org/10.1016/j.ypmed.2021.106790) (2021).
27. Lee, B., Yeon, P. & Park, S. The factors and relationships influencing forest hiking exercise characteristics after COVID-19 occurrence: At Seoul metropolitan area and in hikers' 20s and 30s. *Int. J. Environ. Res. Public Heal.* **19**, 16403, DOI: [10.3390/ijerph192416403](https://doi.org/10.3390/ijerph192416403) (2022).
28. Veitch, J. *et al.* The REVAMP natural experiment study: the impact of a play-scape installation on park visitation and park-based physical activity. *Int. J. Behav. Nutr. Phys. Activity* **15**, DOI: [10.1186/s12966-017-0625-5](https://doi.org/10.1186/s12966-017-0625-5) (2018).
29. Puhakka, R. *et al.* Greening of daycare yards with biodiverse materials affords well-being, play and environmental relationships. *Int. J. Environ. Res. Public Heal.* **16**, 2948, DOI: [10.3390/ijerph16162948](https://doi.org/10.3390/ijerph16162948) (2019).
30. Zang, P. *et al.* Eye-level street greenery and walking behaviors of older adults. *Int. J. Environ. Res. Public Heal.* **17**, 6130, DOI: [10.3390/ijerph17176130](https://doi.org/10.3390/ijerph17176130) (2020).
31. Petrunoff, N. A. *et al.* Associations of park features with park use and park-based physical activity in an urban environment in asia: A cross-sectional study. *Heal. & Place* **75**, 102790, DOI: [10.1016/j.healthplace.2022.102790](https://doi.org/10.1016/j.healthplace.2022.102790) (2022).
32. Krenn, P. J., Oja, P. & Titze, S. Route choices of transport bicyclists: a comparison of actually used and shortest routes. *Int. J. Behav. Nutr. Phys. Activity* **11**, DOI: [10.1186/1479-5868-11-31](https://doi.org/10.1186/1479-5868-11-31) (2014).
33. Lu, Y., Sarkar, C. & Xiao, Y. The effect of street-level greenery on walking behavior: Evidence from hong kong. *Soc. Sci. & Medicine* **208**, 41–49, DOI: [10.1016/j.socscimed.2018.05.022](https://doi.org/10.1016/j.socscimed.2018.05.022) (2018).
34. Zang, P. *et al.* Nonlinear effects of the built environment on light physical activity among older adults: The case of lanzhou, china. *Int. J. Environ. Res. Public Heal.* **19**, 8848, DOI: [10.3390/ijerph19148848](https://doi.org/10.3390/ijerph19148848) (2022).
35. Uijtdewilligen, L. *et al.* The park prescription study: Development of a community-based physical activity intervention for a multi-ethnic asian population. *PLOS ONE* **14**, e0218247, DOI: [10.1371/journal.pone.0218247](https://doi.org/10.1371/journal.pone.0218247) (2019).
36. Liu, Y., Wang, X., Zhou, S. & Wu, W. The association between spatial access to physical activity facilities within home and workplace neighborhoods and time spent on physical activities: evidence from guangzhou, china. *Int. J. Heal. Geogr.* **19**, DOI: [10.1186/s12942-020-00216-2](https://doi.org/10.1186/s12942-020-00216-2) (2020).
37. Müller-Riemenschneider, F. *et al.* Prescribing physical activity in parks to improve health and wellbeing: Protocol of the park prescription randomized controlled trial. *Int. J. Environ. Res. Public Heal.* **15**, 1154, DOI: [10.3390/ijerph15061154](https://doi.org/10.3390/ijerph15061154) (2018).
38. Moran, M. *et al.* Understanding the relationships between the physical environment and physical activity in older adults: a systematic review of qualitative studies. *Int. J. Behav. Nutr. Phys. Activity* **11**, DOI: [10.1186/1479-5868-11-79](https://doi.org/10.1186/1479-5868-11-79) (2014).
39. Zhang, Y. *et al.* The impact of interventions in the built environment on physical activity levels: a systematic umbrella review. *Int. J. Behav. Nutr. Phys. Activity* **19**, DOI: [10.1186/s12966-022-01399-6](https://doi.org/10.1186/s12966-022-01399-6) (2022).
40. Mueller, W. *et al.* Neighbourhood and path-based greenspace in three european countries: associations with objective physical activity. *BMC Public Heal.* **21**, DOI: [10.1186/s12889-021-10259-0](https://doi.org/10.1186/s12889-021-10259-0) (2021).
41. Bonaccorsi, G. *et al.* Impact of the built environment and the neighborhood in promoting the physical activity and the healthy aging in older people: An umbrella review. *Int. J. Environ. Res. Public Heal.* **17**, 6127, DOI: [10.3390/ijerph17176127](https://doi.org/10.3390/ijerph17176127) (2020).
42. Yang, Y., Lu, Y., Yang, L., Gou, Z. & Liu, Y. Urban greenery cushions the decrease in leisure-time physical activity during the COVID-19 pandemic: A natural experimental study. *Urban For. & Urban Green.* **62**, 127136, DOI: [10.1016/j.ufug.2021.127136](https://doi.org/10.1016/j.ufug.2021.127136) (2021).
43. Yi, H., Ng, S. T., Chang, C. M., Low, C. X. E. & Tan, C. S. Effects of neighborhood features on healthy aging in place: the composition and context of urban parks and traditional local coffeeshops in singapore. *BMC Geriatr.* **22**, DOI: [10.1186/s12877-022-03679-z](https://doi.org/10.1186/s12877-022-03679-z) (2022).

44. Petrunoff, N. *et al.* Activity in nature mediates a park prescription intervention's effects on physical activity, park use and quality of life: a mixed-methods process evaluation. *BMC Public Heal.* **21**, DOI: [10.1186/s12889-021-10177-1](https://doi.org/10.1186/s12889-021-10177-1) (2021).
45. Cassarino, M., Bantry-White, E. & Setti, A. Cognitive and sensory dimensions of older people's preferences of outdoor spaces for walking: A survey study in Ireland. *Int. J. Environ. Res. Public Heal.* **16**, 1340, DOI: [10.3390/ijerph16081340](https://doi.org/10.3390/ijerph16081340) (2019).
46. Andersson, D., Wahlgren, L. & Schantz, P. Pedestrians' perceptions of route environments in relation to deterring or facilitating walking. *Front. Public Heal.* **10**, DOI: [10.3389/fpubh.2022.1012222](https://doi.org/10.3389/fpubh.2022.1012222) (2023).
47. McCormack, G. R. & Shiell, A. In search of causality: a systematic review of the relationship between the built environment and physical activity among adults. *Int. J. Behav. Nutr. Phys. Activity* **8**, 125, DOI: [10.1186/1479-5868-8-125](https://doi.org/10.1186/1479-5868-8-125) (2011).
48. Benton, J. S. *et al.* A natural experimental study of improvements along an urban canal: impact on canal usage, physical activity and other wellbeing behaviours. *Int. J. Behav. Nutr. Phys. Activity* **18**, DOI: [10.1186/s12966-021-01088-w](https://doi.org/10.1186/s12966-021-01088-w) (2021).
49. Wang, H., Dai, X., Wu, J., Wu, X. & Nie, X. Influence of urban green open space on residents' physical activity in China. *BMC Public Heal.* **19**, 1–12, DOI: [10.1186/s12889-019-7416-7](https://doi.org/10.1186/s12889-019-7416-7) (2019).
50. Ponjoan, A. *et al.* Impact of residential greenness on myocardial infarction in the population with diabetes: A sex-dependent association? *Environ. Res.* **205**, 112449, DOI: [10.1016/j.envres.2021.112449](https://doi.org/10.1016/j.envres.2021.112449) (2022).
51. Han, H., Yang, K., Yang, C., Yang, G. & Xu, L. Influence and mechanism of a multi-scale built environment on the leisure activities of the elderly: Evidence from Hefei city in China. *Int. J. Environ. Res. Public Heal.* **19**, 9237, DOI: [10.3390/ijerph19159237](https://doi.org/10.3390/ijerph19159237) (2022).
52. Lovasi, G. S., Grady, S. & Rundle, A. Steps forward: Review and recommendations for research on walkability, physical activity and cardiovascular health. *Public Heal. Rev.* **33**, 484–506, DOI: [10.1007/bf03391647](https://doi.org/10.1007/bf03391647) (2011).
53. Richardson, E. A., Pearce, J., Mitchell, R. & Kingham, S. Role of physical activity in the relationship between urban green space and health. *Public Heal.* **127**, 318–324, DOI: [10.1016/j.puhe.2013.01.004](https://doi.org/10.1016/j.puhe.2013.01.004) (2013).
54. Müller-Riemenschneider, F. *et al.* Effectiveness of prescribing physical activity in parks to improve health and wellbeing - the park prescription randomized controlled trial. *Int. J. Behav. Nutr. Phys. Activity* **17**, DOI: [10.1186/s12966-020-00941-8](https://doi.org/10.1186/s12966-020-00941-8) (2020).
55. Ugolini, F. *et al.* Effects of the COVID-19 pandemic on the use and perceptions of urban green space: An international exploratory study. *Urban For. & Urban Green.* **56**, 126888, DOI: [10.1016/j.ufug.2020.126888](https://doi.org/10.1016/j.ufug.2020.126888) (2020).
56. Bohnert, A. M., Nicholson, L. M., Mertz, L., Bates, C. R. & Gerstein, D. E. Green schoolyard renovations in low-income urban neighborhoods: Benefits to students, schools, and the surrounding community. *Am. J. Community Psychol.* **69**, 463–473, DOI: [10.1002/ajcp.12559](https://doi.org/10.1002/ajcp.12559) (2021).
57. Ulmer, J. M. *et al.* Multiple health benefits of urban tree canopy: The mounting evidence for a green prescription. *Heal. & Place* **42**, 54–62, DOI: [10.1016/j.healthplace.2016.08.011](https://doi.org/10.1016/j.healthplace.2016.08.011) (2016).
58. Neale, C. *et al.* The aging urban brain: Analyzing outdoor physical activity using the emotiv affectiv suite in older people. *J. Urban Heal.* **94**, 869–880, DOI: [10.1007/s11524-017-0191-9](https://doi.org/10.1007/s11524-017-0191-9) (2017).
59. Sun, Y., Tan, S., He, Q. & Shen, J. Influence mechanisms of community sports parks to enhance social interaction: A Bayesian belief network analysis. *Int. J. Environ. Res. Public Heal.* **19**, 1466, DOI: [10.3390/ijerph19031466](https://doi.org/10.3390/ijerph19031466) (2022).
60. Tabrizi, N., Lak, A. & Moussavi, A. S. M. R. Green space and the health of the older adult during pandemics: a narrative review on the experience of COVID-19. *Front. Public Heal.* **11**, DOI: [10.3389/fpubh.2023.1218091](https://doi.org/10.3389/fpubh.2023.1218091) (2023).
61. Dennis, M., Cook, P. A., James, P., Wheeler, C. P. & Lindley, S. J. Relationships between health outcomes in older populations and urban green infrastructure size, quality and proximity. *BMC Public Heal.* **20**, DOI: [10.1186/s12889-020-08762-x](https://doi.org/10.1186/s12889-020-08762-x) (2020).
62. Olszewska-Guizzo, A., Sia, A., Fogel, A. & Ho, R. Can exposure to certain urban green spaces trigger frontal alpha asymmetry in the brain?—preliminary findings from a passive task EEG study. *Int. J. Environ. Res. Public Heal.* **17**, 394, DOI: [10.3390/ijerph17020394](https://doi.org/10.3390/ijerph17020394) (2020).
63. Olszewska-Guizzo, A., Sia, A., Fogel, A. & Ho, R. Features of urban green spaces associated with positive emotions, mindfulness and relaxation. *Sci. Reports* **12**, DOI: [10.1038/s41598-022-24637-0](https://doi.org/10.1038/s41598-022-24637-0) (2022).

64. Carver, A., Lorenzon, A., Veitch, J., Macleod, A. & Sugiyama, T. Is greenery associated with mental health among residents of aged care facilities? a systematic search and narrative review. *Aging & Mental Heal.* **24**, 1–7, DOI: [10.1080/13607863.2018.1516193](https://doi.org/10.1080/13607863.2018.1516193) (2018).
65. Sugiyama, T., Carver, A., Sugiyama, M., Lorenzon, A. & Davison, T. E. Views of greenery and psychological well-being in residential aged care facilities: Longitudinal associations. *HERD: Heal. Environ. Res. & Des. J.* **15**, 219–232, DOI: [10.1177/19375867211059757](https://doi.org/10.1177/19375867211059757) (2021).
66. Tran, I., Sabol, O. & Mote, J. The relationship between greenspace exposure and psychopathology symptoms: A systematic review. *Biol. Psychiatry Glob. Open Sci.* **2**, 206–222, DOI: [10.1016/j.bpsgos.2022.01.004](https://doi.org/10.1016/j.bpsgos.2022.01.004) (2022).
67. Olszewska-Guizzo, A. *et al.* Therapeutic garden with contemplative features induces desirable changes in mood and brain activity in depressed adults. *Front. Psychiatry* **13**, DOI: [10.3389/fpsyt.2022.757056](https://doi.org/10.3389/fpsyt.2022.757056) (2022).
68. Moreira, T. C. L. *et al.* Assessing the impact of urban environment and green infrastructure on mental health: results from the são paulo megacity mental health survey. *J. Expo. Sci. & Environ. Epidemiol.* **32**, 205–212, DOI: [10.1038/s41370-021-00349-x](https://doi.org/10.1038/s41370-021-00349-x) (2021).
69. Zamora, A. N. *et al.* Exploring the beliefs and perceptions of spending time in nature among U.S. youth. *BMC Public Heal.* **21**, DOI: [10.1186/s12889-021-11622-x](https://doi.org/10.1186/s12889-021-11622-x) (2021).
70. Helbich, M., O'Connor, R. C., Nieuwenhuijsen, M. & Hagedoorn, P. Greenery exposure and suicide mortality later in life: A longitudinal register-based case-control study. *Environ. Int.* **143**, 105982, DOI: [10.1016/j.envint.2020.105982](https://doi.org/10.1016/j.envint.2020.105982) (2020).
71. Helbich, M., de Beurs, D., Kwan, M.-P., O'Connor, R. C. & Groenewegen, P. P. Natural environments and suicide mortality in the netherlands: a cross-sectional, ecological study. *The Lancet Planet. Heal.* **2**, e134–e139, DOI: [10.1016/s2542-5196\(18\)30033-0](https://doi.org/10.1016/s2542-5196(18)30033-0) (2018).
72. Zeng, C., Lin, W. & Chen, Q. Psychophysiological responses of college students to audio-visual forest trail landscapes. *J. Urban Heal.* **100**, 711–724, DOI: [10.1007/s11524-023-00757-4](https://doi.org/10.1007/s11524-023-00757-4) (2023).
73. Groenewegen, P. P., van den Berg, A. E., de Vries, S. & Verheij, R. A. Vitamin G: effects of green space on health, well-being, and social safety. *BMC Public Heal.* **6**, DOI: [10.1186/1471-2458-6-149](https://doi.org/10.1186/1471-2458-6-149) (2006).
74. Sachs, A. L. *et al.* Rationale, feasibility, and acceptability of the meeting in nature together (MINT) program: A novel nature-based social intervention for loneliness reduction with teen parents and their peers. *Int. J. Environ. Res. Public Heal.* **19**, 11059, DOI: [10.3390/ijerph191711059](https://doi.org/10.3390/ijerph191711059) (2022).
75. Keralis, J. M. *et al.* Health and the built environment in united states cities: measuring associations using google street view-derived indicators of the built environment. *BMC Public Heal.* **20**, DOI: [10.1186/s12889-020-8300-1](https://doi.org/10.1186/s12889-020-8300-1) (2020).
76. Gu, J., Liu, H. & Lu, H. Can even a small amount of greenery be helpful in reducing stress? a systematic review. *Int. J. Environ. Res. Public Heal.* **19**, 9778, DOI: [10.3390/ijerph19169778](https://doi.org/10.3390/ijerph19169778) (2022).
77. Zhang, R., Zhang, C.-Q. & Rhodes, R. E. The pathways linking objectively-measured greenspace exposure and mental health: A systematic review of observational studies. *Environ. Res.* **198**, 111233, DOI: [10.1016/j.envres.2021.111233](https://doi.org/10.1016/j.envres.2021.111233) (2021).
78. Diana, Emanuel, Keren & Izhak. Ethnic differences in home-related maternal stress: Muslim and jewish mothers. *Int. J. Environ. Res. Public Heal.* **16**, 4393, DOI: [10.3390/ijerph16224393](https://doi.org/10.3390/ijerph16224393) (2019).
79. Dzhambov, A. M. *et al.* Multiple pathways link urban green- and bluespace to mental health in young adults. *Environ. Res.* **166**, 223–233, DOI: [10.1016/j.envres.2018.06.004](https://doi.org/10.1016/j.envres.2018.06.004) (2018).
80. Agyemang, C. *et al.* Ethnic differences in the effect of environmental stressors on blood pressure and hypertension in the netherlands. *BMC Public Heal.* **7**, DOI: [10.1186/1471-2458-7-118](https://doi.org/10.1186/1471-2458-7-118) (2007).
81. Zanini, M. J. *et al.* Urban-related environmental exposures during pregnancy and placental development and preeclampsia: a review. *Curr. Hypertens. Reports* **22**, DOI: [10.1007/s11906-020-01088-4](https://doi.org/10.1007/s11906-020-01088-4) (2020).
82. Mueller, W., Milner, J., Loh, M., Vardoulakis, S. & Wilkinson, P. Exposure to urban greenspace and pathways to respiratory health: An exploratory systematic review. *Sci. The Total. Environ.* **829**, 154447, DOI: [10.1016/j.scitotenv.2022.154447](https://doi.org/10.1016/j.scitotenv.2022.154447) (2022).
83. Lv, H. & Wang, R. Association between the built environment and moderate to vigorous leisure-time physical activity among suzhou adolescents: a cross-sectional study. *BMC Public Heal.* **23**, DOI: [10.1186/s12889-023-16243-0](https://doi.org/10.1186/s12889-023-16243-0) (2023).
84. Van Cauwenberg, J., Nathan, A., Barnett, A., Barnett, D. W. & Cerin, E. Relationships between neighbourhood physical environmental attributes and older adults' leisure-time physical activity: A systematic review and meta-analysis. *Sports Medicine* **48**, 1635–1660, DOI: [10.1007/s40279-018-0917-1](https://doi.org/10.1007/s40279-018-0917-1) (2018).

85. Groenewegen, P. P. *et al.* Neighbourhood social and physical environment and general practitioner assessed morbidity. *Heal. & Place* **49**, 68–84, DOI: [10.1016/j.healthplace.2017.11.006](https://doi.org/10.1016/j.healthplace.2017.11.006) (2018).
86. Astell-Burt, T., Walsan, R., Davis, W. & Feng, X. What types of green space disrupt a lonelygenic environment? a cohort study. *Soc. Psychiatry Psychiatr. Epidemiol.* **58**, 745–755, DOI: [10.1007/s00127-022-02381-0](https://doi.org/10.1007/s00127-022-02381-0) (2022).
87. Russo, A. & Cirella, G. Modern compact cities: How much greenery do we need? *Int. J. Environ. Res. Public Heal.* **15**, 2180, DOI: [10.3390/ijerph15102180](https://doi.org/10.3390/ijerph15102180) (2018).
88. Wang, R. *et al.* Urban greenery and mental wellbeing in adults: Cross-sectional mediation analyses on multiple pathways across different greenery measures. *Environ. Res.* **176**, 108535, DOI: [10.1016/j.envres.2019.108535](https://doi.org/10.1016/j.envres.2019.108535) (2019).
89. Orr, N., Wagstaffe, A., Briscoe, S. & Garside, R. How do older people describe their sensory experiences of the natural world? a systematic review of the qualitative evidence. *BMC Geriatr.* **16**, DOI: [10.1186/s12877-016-0288-0](https://doi.org/10.1186/s12877-016-0288-0) (2016).
90. Saadi, D., Schnell, I. & Tirosh, E. Ethnic differences in environmental restoration: Arab and jewish women in israel. *Int. J. Environ. Res. Public Heal.* **18**, 12628, DOI: [10.3390/ijerph182312628](https://doi.org/10.3390/ijerph182312628) (2021).
91. Ng, K. *et al.* Effects of horticultural therapy on asian older adults: A randomized controlled trial. *Int. J. Environ. Res. Public Heal.* **15**, 1705, DOI: [10.3390/ijerph15081705](https://doi.org/10.3390/ijerph15081705) (2018).
92. Schram-Bijkerk, D., Otte, P., Dirven, L. & Breure, A. M. Indicators to support healthy urban gardening in urban management. *Sci. The Total. Environ.* **621**, 863–871, DOI: [10.1016/j.scitotenv.2017.11.160](https://doi.org/10.1016/j.scitotenv.2017.11.160) (2018).
93. Sia, A. *et al.* Nature-based activities improve the well-being of older adults. *Sci. Reports* **10**, DOI: [10.1038/s41598-020-74828-w](https://doi.org/10.1038/s41598-020-74828-w) (2020).
94. Mirzaei, M., Verrelst, J., Arbabi, M., Shaklabadi, Z. & Lotfizadeh, M. Urban heat island monitoring and impacts on citizen's general health status in isfahan metropolis: A remote sensing and field survey approach. *Remote. Sens.* **12**, 1350, DOI: [10.3390/rs12081350](https://doi.org/10.3390/rs12081350) (2020).
95. Kumar, P. *et al.* The nexus between air pollution, green infrastructure and human health. *Environ. Int.* **133**, 105181, DOI: [10.1016/j.envint.2019.105181](https://doi.org/10.1016/j.envint.2019.105181) (2019).
96. Zutter, C. & Stoltz, A. Community gardens and urban agriculture: Healthy environment/healthy citizens. *Int. J. Mental Heal. Nurs.* DOI: [10.1111/inm.13149](https://doi.org/10.1111/inm.13149) (2023).
97. Wong, G. C. L. *et al.* Horticultural therapy reduces biomarkers of immunosenescence and inflammaging in community-dwelling older adults: A feasibility pilot randomized controlled trial. *The Journals Gerontol. Ser. A* **76**, 307–317, DOI: [10.1093/gerona/glaa271](https://doi.org/10.1093/gerona/glaa271) (2020).
98. Benton, J. S. *et al.* Evaluating the impact of improvements in urban green space on older adults' physical activity and wellbeing: protocol for a natural experimental study. *BMC Public Heal.* **18**, DOI: [10.1186/s12889-018-5812-z](https://doi.org/10.1186/s12889-018-5812-z) (2018).
99. da Cunha, M. A., Paraguassú, L. A. A., de Aquino Assis, J. G., de Paula Carvalho Silva, A. B. & de Cassia Vieira Cardoso, R. Urban gardening and neglected and underutilized species in salvador, bahia, brazil. *J. Ethnobiol. Ethnomedicine* **16**, DOI: [10.1186/s13002-020-00421-0](https://doi.org/10.1186/s13002-020-00421-0) (2020).
100. van den Bosch, M. A. & Depledge, M. H. Healthy people with nature in mind. *BMC Public Heal.* **15**, DOI: [10.1186/s12889-015-2574-8](https://doi.org/10.1186/s12889-015-2574-8) (2015).
101. Chiumento, A. *et al.* A haven of green space: learning from a pilot pre-post evaluation of a school-based social and therapeutic horticulture intervention with children. *BMC Public Heal.* **18**, DOI: [10.1186/s12889-018-5661-9](https://doi.org/10.1186/s12889-018-5661-9) (2018).
102. Paddle, E. & Gilliland, J. Orange is the new green: Exploring the restorative capacity of seasonal foliage in schoolyard trees. *Int. J. Environ. Res. Public Heal.* **13**, 497, DOI: [10.3390/ijerph13050497](https://doi.org/10.3390/ijerph13050497) (2016).
103. Kwon, O.-H. *et al.* Urban green space and happiness in developed countries. *EPJ Data Sci.* **10**, DOI: [10.1140/epjds/s13688-021-00278-7](https://doi.org/10.1140/epjds/s13688-021-00278-7) (2021).
104. Zhou, Y., Yuan, Y., Chen, Y. & Lai, S. Association pathways between neighborhood greenspaces and the physical and mental health of older adults—a cross-sectional study in guangzhou, china. *Front. Public Heal.* **8**, DOI: [10.3389/fpubh.2020.551453](https://doi.org/10.3389/fpubh.2020.551453) (2020).
105. Ruijsbroek, A. *et al.* Neighbourhood green space, social environment and mental health: an examination in four european cities. *Int. J. Public Heal.* **62**, 657–667, DOI: [10.1007/s00038-017-0963-8](https://doi.org/10.1007/s00038-017-0963-8) (2017).
106. Jennings, V. & Bamkole, O. The relationship between social cohesion and urban green space: An avenue for health promotion. *Int. J. Environ. Res. Public Heal.* **16**, 452, DOI: [10.3390/ijerph16030452](https://doi.org/10.3390/ijerph16030452) (2019).

107. Hong, A. *et al.* Linking green space to neighborhood social capital in older adults: The role of perceived safety. *Soc. Sci. & Medicine* **207**, 38–45, DOI: [10.1016/j.socscimed.2018.04.051](https://doi.org/10.1016/j.socscimed.2018.04.051) (2018).
108. Maas, J., Van Dillen, S. M., Verheij, R. A. & Groenewegen, P. P. Social contacts as a possible mechanism behind the relation between green space and health. *Heal. & place* **15**, 586–595, DOI: [10.1016/j.healthplace.2008.09.006](https://doi.org/10.1016/j.healthplace.2008.09.006) (2009).
109. Rostami, R., Lamit, H., Khoshnava, S. M. & Rostami, R. The role of historical persian gardens on the health status of contemporary urban residents. *EcoHealth* **11**, 308–321, DOI: [10.1007/s10393-014-0939-6](https://doi.org/10.1007/s10393-014-0939-6) (2014).
110. Du, Y., Jiang, H., Huang, Z. & Yang, H. Associations between neighborhood environment and sense of community belonging in urban china: Examining mediation effects of neighborly interactions and community satisfaction. *Front. Public Heal.* **10**, DOI: [10.3389/fpubh.2022.1105473](https://doi.org/10.3389/fpubh.2022.1105473) (2023).
111. Dadvand, P. *et al.* Use of green spaces, self-satisfaction and social contacts in adolescents: A population-based CASPIAN-v study. *Environ. Res.* **168**, 171–177, DOI: [10.1016/j.envres.2018.09.033](https://doi.org/10.1016/j.envres.2018.09.033) (2019).
112. Włodarczyk-Marciniak, R., Sikorska, D. & Krauze, K. Residents' awareness of the role of informal green spaces in a post-industrial city, with a focus on regulating services and urban adaptation potential. *Sustain. Cities Soc.* **59**, 102236, DOI: [10.1016/j.scs.2020.102236](https://doi.org/10.1016/j.scs.2020.102236) (2020).
113. Jennings, V., Larson, L. & Yun, J. Advancing sustainability through urban green space: Cultural ecosystem services, equity, and social determinants of health. *Int. J. Environ. Res. Public Heal.* **13**, 196, DOI: [10.3390/ijerph13020196](https://doi.org/10.3390/ijerph13020196) (2016).
114. Pinto, L., Ferreira, C. S. & Pereira, P. Environmental and socioeconomic factors influencing the use of urban green spaces in coimbra (portugal). *Sci. The Total. Environ.* **792**, 148293, DOI: [10.1016/j.scitotenv.2021.148293](https://doi.org/10.1016/j.scitotenv.2021.148293) (2021).
115. Nakau, M. *et al.* Spiritual care of cancer patients by integrated medicine in urban green space: A pilot study. *EXPLORE* **9**, 87–90, DOI: [10.1016/j.explore.2012.12.002](https://doi.org/10.1016/j.explore.2012.12.002) (2013).
116. Rosa, C. D., Larson, L. R., Collado, S. & Profice, C. C. Forest therapy can prevent and treat depression: Evidence from meta-analyses. *Urban For. & Urban Green.* **57**, 126943, DOI: [10.1016/j.ufug.2020.126943](https://doi.org/10.1016/j.ufug.2020.126943) (2021).
117. Quercia, D., Lathia, N., Calabrese, F., Di Lorenzo, G. & Crowcroft, J. Recommending social events from mobile phone location data. In *IEEE International Conference on Data Mining*, 971–976, DOI: [10.1109/icdm.2010.152](https://doi.org/10.1109/icdm.2010.152) (IEEE, 2010).
118. Hu, Y., Koren, Y. & Volinsky, C. Collaborative filtering for implicit feedback datasets. In *IEEE International Conference on Data Mining*, 263–272, DOI: [10.1109/icdm.2008.22](https://doi.org/10.1109/icdm.2008.22) (IEEE, 2008).
119. Brown, T. B. *et al.* Language models are few-shot learners, DOI: [10.48550/arxiv.2005.14165](https://doi.org/10.48550/arxiv.2005.14165) (2020). [2005.14165](https://arxiv.org/abs/2005.14165).
120. Achiam, J. *et al.* GPT-4 technical report, DOI: [10.48550/arxiv.2303.08774](https://doi.org/10.48550/arxiv.2303.08774) (2023). [2303.08774](https://arxiv.org/abs/2303.08774).
121. OpenAI. OpenAI API Reference. <https://platform.openai.com/docs/api-reference/introduction> (2024). [Online; accessed 12-July-2024].
122. OpenAI. Prompt engineering – six strategies for getting better results. <https://platform.openai.com/docs/guides/prompt-engineering> (2024). [Online; accessed 12-July-2024].
123. OpenStreetMap Wiki. Map features. https://wiki.openstreetmap.org/w/index.php?title=Map_features&oldid=2543361 (2024). [Online; accessed 12-July-2024].
124. Li, L.-J., Socher, R. & Fei-Fei, L. Towards total scene understanding: Classification, annotation and segmentation in an automatic framework. In *Conference on Computer Vision and Pattern Recognition*, 2036–2043, DOI: [10.1109/cvpr.2009.5206718](https://doi.org/10.1109/cvpr.2009.5206718) (IEEE, 2009).
125. Thomee, B. *et al.* YFCC100M: The new data in multimedia research. *Commun. ACM* **59**, 64–73, DOI: [10.1145/2812802](https://doi.org/10.1145/2812802) (2016).
126. Reimers, N. & Gurevych, I. Sentence-BERT: Sentence embeddings using siamese BERT-networks. In *Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing*, DOI: [10.18653/v1/d19-1410](https://doi.org/10.18653/v1/d19-1410) (ACL, 2019).
127. Lugaresi, C. *et al.* Mediapipe: A framework for building perception pipelines, DOI: [10.48550/ARXIV.1906.08172](https://doi.org/10.48550/ARXIV.1906.08172) (2019).
128. MediaPipe Authors. Language detection guide for Python. https://developers.google.com/mediapipe/solutions/text/language_detector/python (2023). [Online; accessed 12-July-2024].
129. Tiedemann, J. & Thottingal, S. OPUS-MT – Building open translation services for the World. In *22nd Annual Conference of the European Association for Machine Translation* (Lisbon, Portugal, 2020).
130. Miles, M. B. & Huberman, A. M. *Qualitative data analysis: An expanded sourcebook* (Sage, 1994).
131. Braun, V. & Clarke, V. Using thematic analysis in psychology. *Qual. research psychology* **3**, 77–101, DOI: [10.1191/1478088706qp063oa](https://doi.org/10.1191/1478088706qp063oa) (2006).